

Ronald E. Jung  
 Anja Zembic  
 Bjarni E. Pjetursson  
 Marcel Zwahlen  
 Daniel S. Thoma

# Systematic review of the survival rate and the incidence of biological, technical, and aesthetic complications of single crowns on implants reported in longitudinal studies with a mean follow-up of 5 years

## Authors' affiliations:

Ronald E. Jung, Daniel S. Thoma, Clinic of Fixed and Removable Prosthodontics and Dental Material Science, University of Zurich, Zurich, Switzerland  
 Anja Zembic, Implantology Department ACTA, Amsterdam, the Netherlands  
 Bjarni E. Pjetursson, Reconstructive Dentistry, Faculty of Odontology, University of Iceland, Reykjavik, Iceland  
 Marcel Zwahlen, Institute of Social and Preventive Medicine, University of Berne, Berne, Switzerland

## Corresponding author:

Ronald E. Jung  
 Clinic of Fixed and Removable Prosthodontics and Dental Material Science  
 University of Zurich  
 Plattenstrasse 11  
 CH-8032 Zurich, Switzerland  
 Tel.: +41 44 634 3251  
 Fax: +41 44 634 4305  
 e-mail: ronald.jung@zzm.uzh.ch

## Conflicts of interest:

The authors report no conflicts of interest.

**Key words:** crown dental implants, humans, survival in humans, systematic review

## Abstract

**Objective:** To assess the 5-year survival of implant-supported single crowns (SCs) and to describe the incidence of biological, technical, and aesthetic complications. The focused question was: What is the survival rate of implants supporting single crowns and implant-supported crowns with a mean follow-up of 5 years and to which extent do biological, technical, and aesthetic complications occur?

**Methods:** A Medline search (2006–2011) was performed for clinical studies focusing on implant-supported SCs with a mean follow-up of at least 5 years. The search was complemented by an additional hand search and the inclusion of 24 studies from a previous systematic review (Jung et al. 2008a). Survival and complication rates were analyzed using random-effects Poisson's regression models to obtain summary estimates of 5- and 10-year proportions.

**Results:** Forty-six studies derived from an initial search count of 1083 titles and the complementary publications from the previous systematic review (Jung et al. 2008a) were selected and the data were extracted. Based on the meta-analysis, survival of implants supporting SCs at 5 years amounted to 97.2% (95% CI: 96.3–97.9%), and at 10 years amounted to 95.2% (95% CI: 91.8–97.2%). The survival of implant-supported SCs was 96.3% (95% CI: 94.2–97.6%) after 5 years and 89.4% (95% CI: 82.8–93.6%) after 10 years. For biological complications, a 5-year cumulative soft tissue complication rate of 7.1% (95% CI: 4.4–11.3%) and a cumulative complication rate for implants with bone loss >2 mm of 5.2% (95% CI: 3.1–8.6%) were calculated. Technical complications reached a cumulative incidence of 8.8% (95% CI: 5.1–15.0%) for screw-loosening, 4.1% (95% CI: 2.2–7.5%) for loss of retention, and 3.5% (95% CI: 2.4–5.2%) for fracture of the veneering material after 5 years. The cumulative 5-year aesthetic complication rate amounted to 7.1% (95% CI: 3.6–13.6%).

**Conclusions:** The outcomes of the meta-analysis demonstrated high implant survival rates for both the single tooth implants and the respective single crowns after 5 and 10 years. However, technical, biological, and aesthetic complications were frequent.

## Date:

Accepted 09 June 2012

## To cite this article:

Jung RE, Zembic A, Pjetursson BE, Zwahlen M, Thoma DS. Systematic review of the survival rate and the incidence of biological, technical and esthetic complications of single crowns on implants reported in longitudinal studies with a mean follow-up of 5 years. *Clin. Oral Implants Res.* 23(Suppl. 6), 2012, 2–21  
 doi: 10.1111/j.1600-0501.2012.02547.x

The advent of osseointegration has had a fundamental impact on the therapeutic approaches and strategies implemented today in the field of prosthetic rehabilitation of various types of edentulism. The results from better oral prophylaxis and maintenance of patients have led today to a shift from fully edentulous patients to an increased number

of partially edentulous jaws and single tooth gaps. Hence, the treatment of a single tooth gap worldwide has become an important indication within the daily dental practice.

When it comes to the decision-making process between implant-supported single crowns (SCs) and tooth-supported fixed dental prosthesis (FDP), the related decision cri-

teria should be essentially derived from scientific evidence and objective surgically/prosthetically oriented risk assessments as well as patient-related factors including cost effectiveness and quality of life. In terms of a hierarchy of decisions the most important question is, whether or not the prognosis of implant-supported reconstruction is similar to those of tooth-supported FDP.

To answer this question on the highest level of evidence, the use of systematic reviews has been proposed to be an appropriate method (Egger et al. 2001). Hence, systematic reviews are employed in medicine and dentistry to summarize cumulative information on the optimal treatment for clinically important questions. Based on the results of systematic reviews, the clinicians should be able to make appropriate decisions and recommendations for individual clinical indications and to treat patients in an evidence-based way.

A former systematic review of the survival and complication rates of implant-supported SCs was performed from the years 1966 to 2006 (Jung et al. 2008a). During this time period 26 prospective and retrospective cohort studies met the inclusion criteria. In a meta-analysis of these studies the survival rates of implant-supporting SCs was 96.8% (95% confidence interval (CI): 95.9–97.6%) after 5 years. The survival rate of SCs supported by implants was 94.5% (95% CI: 92.5–95.9%) after 5 years of function. This information helped the dentists worldwide in their decision-making process and to inform the patients about the treatment outcomes. However, this information is only valuable when it is going to be continuously updated to prevent the clinicians from using the most current data derived from the literature.

Therefore, it was decided to perform an additional literature search from 2006 to 2011 to identify clinical studies reporting on implant-supported SCs and to update the former systematic review (Jung et al. 2008a).

The objective of this systematic review was to assess the 5-year survival of implant-supported SCs and to describe the rate of biological, technical, and aesthetic complications.

## Material and methods

### Search strategy

This systematic review was designed as an update to a previously prepared publication with the same objectives (Jung et al. 2008a). For that purpose, a Medline (PubMed) search was performed for clinical studies,

including articles published from 1 August 2006 up to 31 August 2011 in the dental literature. The search was limited to the English and German language. In addition, full-text articles of reviews published between January 2008 and August 2011 were obtained. An additional hand search was performed identifying relevant studies by screening these reviews and the reference list of all included publications (reference list “list of reviews”)

### Search terms

The following search terms (all MeSH terms) were selected: “dental implants” AND (“crowns” OR “survival”). The search was limited to “humans” (MeSH term), “Dental Journals”, and “Medline”.

### Inclusion criteria

Clinical publications were considered if all the following criteria were suitable: (i) human trials with a minimum amount of 10 patients with SCs; (ii) mean follow-up of at least 5 years in function; (iii) randomized controlled trials (RCT), controlled clinical trials (CCT), prospective case series, cohort studies, and retrospective studies; (iv) patients needed to be examined clinically; and (v) reported details of suprastructures.

### Exclusion criteria

Studies not meeting all inclusion criteria were excluded from the review. Publications dealing with the following topics were also excluded: studies not reporting in detail the prosthodontic component, reports based on questionnaires, interviews, and charts.

### Selection of studies

Two authors (DTH and AZE) independently screened the titles derived from this broad search based on the inclusion criteria. Disagreements were resolved by discussion. Following this, abstracts of all titles agreed on by both authors were obtained and screened for meeting the inclusion criteria. If no abstract was available in the database, the abstract of the printed article was used. Based on the selection of abstracts, articles were then obtained in full text. If title and abstract did not provide sufficient information regarding the inclusion criteria, the full report was obtained as well. Again, disagreements were resolved by discussion and Cohen's Kappa-coefficient was calculated as a measure of agreement between the two readers.

Finally, the selection based on inclusion/exclusion criteria was made for the full-text articles. For this purpose materials and meth-

ods, results, and discussions of these studies were screened. This step was carried out by three readers (RJU, DTH, and AZE) and double-checked. Any questions that came up during the screening were discussed within the group to aim for consensus. In addition, all but two publications (24 studies) from the previous systematic review (Jung et al. 2008a) were included in the analyses.

### Data extraction and method of analysis

Due to the high number of included articles, three reviewers (RJU, DTH, and AZE) extracted the data. For standardization purposes, five of the included studies were randomly selected and data extracted independently by all three readers. Any disagreements were discussed to aim for consensus and to standardize the subsequent analyses. The three reviewers then independently extracted the data of all included studies using data extraction tables. In case the publication did not provide sufficient information, authors of the respective publication were contacted by e-mail. In addition, data of the included publications of the previously published review (Jung et al. 2008a) were extracted as well. All extracted data were double-checked, and any questions that came up during the screening and the data extraction were discussed within the group to aim for consensus.

Information on the following parameters was extracted: author(s), year of publication, implant system, study design, number of patients, number of implants, number of crowns, dropouts, reconstruction material, type of fixation, follow-up, implant and crown survival, as well as the number of complications (technical, biological) and aesthetic outcomes. Based on the included studies, the number of events for all technical, biological and aesthetic complications was extracted and the corresponding total exposure time of the reconstruction was calculated.

### Statistical analysis

By definition, failure and complication rates are calculated by dividing the number of events (failures or complications) in the numerator by the total exposure time (SC-time and/or implant-time) in the denominator.

The numerator could usually be extracted directly from the publication. The total exposure time was calculated by taking the sum of:

1. Exposure time of SCs/implants that could be followed for the whole observation time.

2. Exposure time up to a failure of the SCs/implants that were lost due to failure during the observation time
3. Exposure time up to the end of observation time for SCs/implants that did not complete the observation period due to reasons such as death, change of address, refusal to participate, nonresponse, chronic illnesses, missed appointments, and work commitments.

For each study, event rates for SCs and/or implants were calculated by dividing the total number of events by the total SCs or implant exposure time in years. For additional analysis, the total number of events was considered to be Poisson distributed for a given sum of implant exposure years and Poisson regression with a logarithmic link-function and total exposure time per study as an offset variable were used (Kirkwood & Sterne 2003b, a).

Robust standard errors were calculated to obtain 95% CI of the summary estimates of the event rates. To assess heterogeneity of the study-specific event rates, the Spearman goodness-of-fit statistics and associated *P*-value were calculated. If the goodness-of-fit *P*-value was below 0.05 indicating heterogeneity, random-effects Poisson regression (with Gamma-distributed random-effects) was used to obtain a summary estimate of the event rates. Five-year and 10-year survival proportions were calculated through the relationship between event rate and survival function  $S$ ,  $S(T) = \exp(-T \cdot \text{event rate})$ , by assuming constant event rates (Kirkwood & Sterne 2003b, a). The 95% CI for the survival proportions were calculated by using the 95% confidence limits of the event rates.

Multivariable Poisson regression was used to investigate formally whether event rates varied by reconstruction material (metal abutment plus metal-ceramic vs. all-ceramic reconstructions) or crown design (cemented vs. screw retained).

All analyses were performed using Stata<sup>®</sup>, version 12.0 (StataCorp, College Station, TX, USA).

## Results

### Study characteristics

The electronic search identified a total of 1083 titles (for details refer to Fig. 1). From assessing the titles, 667 were excluded after discussion. The resulting number of abstracts obtained was 416 of which 224 were excluded (inter-reader agreement  $k = 0.88 \pm 0.87$ ). Thereafter, 192 full-text articles were obtained including 36 review articles. Hand searching provided four more studies. Finally, 22 articles




Fig. 1. Search strategy. \*For details see reference lists “List of reviews” and “List of excluded full text articles and the reasons for exclusion”.

met the inclusion criteria. Twenty-four publications were included from the previously published review (Jung et al. 2008a). This resulted in a final number of 46 publications for the present data analyses (Table 1).

### Exclusion of studies

The reasons for excluding studies ( $n = 138$ , see reference list “List of excluded full-text articles and the reason for exclusion”) after the full text was obtained were: less than 10 patients or SCs (six studies), chart review without clinical examination (5), edentulous patient/no SCs (1), mean follow-up less than 5 years (30), mixed data with no information

on SCs (2), multiple publications on same patient cohort (2), no clinical study (1), limited information on failed implants (1), limited information on prosthetics (60), no information on prosthetics (26), not all patients clinically examined (1), only demographic data (1), tooth-supported SCs (1), only provisional crowns (1). Two publications (Buser et al. 1996; Andersen et al. 2002) from the previous review (Jung et al. 2008a) were excluded due to insufficient number of patients.

### Included studies

The 46 studies that met the inclusion criteria are presented in Table 1. All studies were

**Table 1.** Study and patient characteristics of the reviewed studies

Study	Implant System	Study design	Number of patients	Drop-out (%)	Age range	Mean age	Setting
Jung et al. (2012b)	Straumann	Prospective	29	6.9	28–87	53–60	University
Jung et al. (2012a)	Brånemark/3i/IMZ	Retrospective	20	10	32–87	67.5	University
Schneider et al. (2011)	Straumann/Brånemark	Retrospective	70	NR	19.8–76.6	50.7	University
Bonde et al. (2010)	Brånemark	Retrospective	51	5.9	19–79	43	University students
Krennmair et al. (2010)	Camlog	Retrospective	216	8.3	NR	54.3	Private practice
Matarasso et al. (2010)	Brånemark/Straumann	Retrospective	80	0.0	0	47	University
Schmidlin et al. (2010)	Straumann	Retrospective	64	35.9	33–83	60	University
Urdaneta et al. (2010)	Bicon	Retrospective	108	25.0	27.8–91.8	58.7	Specialist clinic
Zafiroopoulos et al. (2010)	Straumann/Camlog	Retrospective	252	4.4	43–70	49	Private practice
Krieger et al. (2009)	Straumann	Retrospective	49	4.1	16.6–24.7	19.3	University
MacDonald et al. (2009)	Endopore	Prospective	20	15.0	NR	43.5	University
Vigolo & Givani (2009)	3i	Prospective	144	0.0	25–55	37	Private practice
Gottfredsen (2009)	Astra Tech	Prospective	20	5.0	18–59	33	University
Degidi et al. (2008)	BioHorizons	Prospective	155	0.0	18–78	54	Private practice
Hälg et al. (2008)	Straumann	Retrospective	54	1.9	25–68	50.2	Private practice
Jemt (2009)	Brånemark	Retrospective	35	31.4	18–72	32	Specialist clinic
Jemt (2008)	Brånemark	Retrospective	38	32	NR	25.4	Specialist clinic
Schropp & Isidor (2008)	3i	Prospective	45	24.4	20–74	48	University
Pikner et al. (2008)	Brånemark	Retrospective	1346	52.5	NR	NR	Specialist clinic
Kreissl et al. (2007)	3i	Prospective	76	0	18–76	45	University
De Boever et al. (2006)	Straumann	Retrospective	105	0.0	25–86	59.1	University
Romeo et al. (2006)	Straumann	Prospective	129	17.8	NR	53	University
Wagenberg & Froum (2006)	Brånemark, 3i	Retrospective	891*	NR	14–94	57.9	Specialist clinic
Bornstein et al. (2005)	ITI	Prospective	28	4	NR	NR	University
Elkhoury et al. (2005)	3i	Retrospective	39	NR	NR	49.2	University
De Boever & De Boever (2005)	ITI	Prospective	16	0	25–61	NR	University
Wennström et al. (2005)	Astra Tech	Prospective	40	9	20–71	40.9	University
Levin et al. (2005)	NR	Retrospective	48	NR	18–65	36.2	Specialist
Jemt & Lekholm (2005)	Brånemark	Prospective	10	20	21–36	26.3	Specialist clinic
Brägger et al. (2005)	ITI	Prospective	48	30	19–78	49.3	University
Taylor et al. (2004)	BioloK	Prospective	39	0	NR	NR	University
Bernard et al. (2004)	ITI	Retrospective	28	NR	15–55	31	University
Romeo et al. (2004)	ITI	Prospective	250*	14	20–67	53	Private practice
Bianchi & Sanfilippo (2004)	ITI	Prospective	116	4	19–73	45.5	University
Gottfredsen (2004)	Astra Tech	Prospective	20	0	18–59	33	University
Haas et al. (2002)	Brånemark	Prospective	71	3	NR	32	University
Gibbard & Zarb (2002)	Brånemark	Prospective	42	8	15–64	33.4	University
Mericske-Stern et al. (2001)	ITI	Prospective	72	0	19–82	50.1	University
Palmer et al. (2000)	Astra Tech	Prospective	15	7	16–48	49.5	University
Vigolo & Givani (2000)	3i	Retrospective	44	0	18–74	35	Specialist clinic
Thilander et al. (1999)	Brånemark	Prospective	10	0	14–19	15.3	Specialist clinic
Polizzi et al. (1999)	Brånemark	Prospective	21	NR	13–58	30	Specialist clinic
Andersson et al. (1998a)	Brånemark	Prospective	38	8	20–45	31	Specialist clinic and private practice
Andersson et al. (1998b)	Brånemark	Prospective	57	9	NR	32	University
Scheller et al. (1998)	Brånemark multicenter, 12 centers	Prospective	82	18	14–73	35	University and private practice
Henry et al. (1996)	Brånemark multicenter, 7 centers	Prospective	92	16	NR	NR	University and private practice

\*Total number of patients in the study with various types of reconstructions. NR, not reported

published between 1996 and 2012. A total of 27 of the studies were prospective, whereas the remaining 19 were retrospective studies (Table 1). The patients were treated at university settings (29 studies), at specialist clinics (11 studies), or in private practices (6 studies). Two of the studies were multicenter studies (Henry et al. 1996; Scheller et al. 1998). A

total number of 3223 implants were placed in patients with age range 13–94 years. The dropout rate varied between 0% and 52.5%, but was not reported in six studies (Table 1).

The studies reported on 10 commercially available implant systems: 3i Implants (Implant Innovations, Palm Beach Gardens, FL, USA), Astra Tech Implants Dental System

(Astra Tech AB, Mölndal, Sweden), Bicon Dental Implants (Bicon, Boston, MA, USA), BioHorizons Dental Implants (BioHorizons, Birmingham, AL, USA), BioloK Implants (BioHorizons, Birmingham, AL, USA), Brånemark System (Nobel Biocare AG, Zurich, Switzerland), CAMLOG (CAMLOG Biotechnologies AG, Stuttgart, Germany), Endopore Dental

Implants (Sybron Implant Solutions), IMZ implants (Dentsply-Friudent, Mannheim, Germany), ITI/Straumann Dental Implant System (Straumann AG, Waldenburg, Switzerland). Only one study did not report on the commercial name of the implant system that had been used (Levin et al. 2005).

The 46 studies included a total number of 3199 SCs. The material of the reconstruction

was reported in 28 studies and included metal-ceramic (76%), gold-resin (14%), or all-ceramic (10%). Thirty percent of the crowns were screw-retained, whereas 70% were cemented (Table 2).

In 26 studies, all patients in the respective cohorts were followed for the same observation period (5, 10, or 15 years), whereas in 20 studies, variable observation periods were

reported with follow-up time-points between 1 and 26 years (Table 2).

#### Implant survival

All 46 studies reported on implant survival rates (Tables 3 and 4). At the beginning of the studies, 3223 implants were placed. Of these, 104 were known to be lost. Forty-one implants were lost before loading (1.3% of all placed

**Table 2.** Information on implants and SCs in the reviewed studies

Study	Number of implants	Number of crowns	Metal/ceramic	Gold/resin	All-ceramic	Cemented	Screw-retained	Follow-up range	Mean follow-up time
Jung et al. (2012b)	29	29	NR	NR	NR	NR	NR	NR	4.7
Jung et al. (2012a)	20	20	NR	NR	NR	NR	NR	12–14	12.5
Schneider et al. (2011)	100	100	100	0	0	74	26	4.7–11.7	6.2
Bonde et al. (2010)	55	52	0	0	52	52	0	7.5–12	9.4
Krennmair et al. (2010)	112	112	NR	NR	NR	NR	NR	5–7	5.7
Matarasso et al. (2010)	80	80	NR	NR	NR	NR	NR	NR	9.7
Schmidlin et al. (2010)	39	39	39	0	0	35	4	0.8–26.4	6.2
Urdaneta et al. (2010)	326	326	82	228	16	0	326	NR	5.9
Zafropoulos et al. (2010)	252	252	252	0	0	252	0	NR	4.8
Krieger et al. (2009)	24	24	24	0	0	NR	NR	4.6–15.3	8.0
MacDonald et al. (2009)	20	20	20	0	0	0	20	7–9	7.7
Vigolo & Givani (2009)	182	182	182	0	0	182	0	NR	5.0
Gotfredsen (2009)	20	20	20	0	0	20	0	NR	10.0
Degidi et al. (2008)	45	45	NR	NR	NR	NR	NR	NR	5.0
Hälg et al. (2008)	22	22	22	0	0	22	0	3–12.7	5.0
Jemt (2009)	41	41	41	0	0	23	18	NR	10.0
Jemt (2008)	47	47	47	0	0	0	47	NR	12.3
Schropp & Isidor (2008)	45	42	42	0	0	40	2	NR	4.7
Pikner et al. (2008)	45	45	NR	NR	NR	NR	NR	NR	5.0
Kreissl et al. (2007)	46	46	46	0	0	0	46	NR	5.0
De Boever et al. (2006)	80	80	NR	NR	NR	NR	NR	3.3–12	5.2
Romeo et al. (2006)	58	58	58	NR	NR	49	9	3–14	5.0
Wagenberg & Froum (2006)	401	383	NR	NR	NR	NR	NR	1–16	5.9
Bornstein et al. (2005)	39	39	NR	NR	NR	NR	NR	5	5
Elkhoury et al. (2005)	39	39	NR	NR	NR	NR	NR	5	5
De Boever & De Boever (2005)	10	10	NR	NR	NR	NR	NR	3–10*	5
Wennström et al. (2005)	45	44	44	0	0	44	0	5	5
Levin et al. (2005)	30	29	NR	NR	NR	NR	NR	3–10*	5.1
Jemt & Lekholm (2005)	10*	10	10	0	0	10	0	5	5
Brägger et al. (2005)	69	69	69	0	0	67	2	8–12	10
Taylor et al. (2004)	39	38	NR	NR	NR	NR	NR	5	5
Bernard et al. (2004)	32	32	32	0	0	NR	NR	2–9	5
Romeo et al. (2004)	123	121	121	0	0	NR	NR	1–7	5.8
Bianchi & Sanfilippo (2004)	116	116	116	0	0	116	0	1–9	5.2
Gotfredsen (2004)	20	20	20	0	0	20	0	5	5
Haas et al. (2002)	76	75	NR	NR	NR	75	0	4–10	5.5
Gibbard & Zarb (2002)	49	48	NR	NR	NR	2	46	4–13	5.9
Mericske-Stern et al. (2001)	26	26	24	0	0	2	24	5–9	6.5
Palmer et al. (2000)	15	15	15	0	0	15	0	5	5
Vigolo & Givani (2000)	52	52	36	16	0	52	0	5	5
Thilander et al. (1999)	15	15	NR	NR	NR	NR	NR	8	5
Polizzi et al. (1999)	30	30	30	0	0	30	0	3–7	5.3
Andersson et al. (1998a)	38	38	NR	NR	NR	NR	NR	5	5
Andersson et al. (1998b)	65	65	3	0	62	65	0	5	5
Scheller et al. (1998)	99	97	16	0	81	97	0	5	5
Henry et al. (1996)	107	106	61	45	0	NR	NR	5	5
<b>Total</b>	<b>3223</b>	<b>3199</b>	<b>1572</b>	<b>289</b>	<b>211</b>	<b>1344</b>	<b>570</b>	<b>1–26.4</b>	<b>6.2</b>

\*Implants with less than 3 years follow-up time were excluded from the meta-analysis. NR, not reported

**Table 3.** Annual failure rates and 5-year survival of implants

Study	Total number of implants	Mean follow-up time	Number of failure	Total implant exposure time	Estimated failure rate (per 100 implant years)	Estimated survival rate after 5 years (in percent)
<b>Prospective studies</b>						
Jung et al. (2012b)	29	4.7	0	137	0	100.0
Vigolo & Givani (2009)	182	5	0	910	0	100.0
Degidi et al. (2008)	45	5	0	225	0	100.0
Schropp & Isidor (2008)	45	4.7	3	210	1.43	93.1
Kreissl et al. (2007)	46	5	1	230	0.43	97.8
Romeo et al. (2006)	58	5	1	288	0.35	98.3
Bornstein et al. (2005)	39	5	0	190	0	100.0
De Boever & de Boever (2005)	10	5	1	50	2	90.5
Wennström et al. (2005)	45	5	1	208	0.48	97.6
Jemt & Lekholm (2005)	10	5	0	48	0	100.0
Taylor et al. (2004)	39	5	1	190	0.53	97.4
Romeo et al. (2004)	123	5.8	7	711	0.98	95.2
Bianchi & Sanfilippo (2004)	116	5.2	0	594	0	100.0
Gotfredsen (2004)	20	5	0	100	0	100.0
Haas et al. (2002)	76	5.5	5	407	1.23	94.0
Gibbard & Zarb (2002)	49	5.9	1	287	0.35	98.3
Mericske-Stern et al. (2001)	26	6.5	2	169	1.18	94.3
Palmer et al. (2000)	15	5	0	70	0	100.0
Polizzi et al. (1999)	30	5.3	1	158	0.63	96.9
Andersson et al. (1998a)	38	5	0	182	0	100.0
Andersson et al. (1998b)	65	5	1	305	0.33	98.4
Scheller et al. (1998)	99	5	3	411	0.73	96.4
Henry et al. (1996)	107	5	3	477	0.63	96.9
<b>Total</b>	<b>1312</b>	<b>5.2</b>	<b>31</b>	<b>6557</b>		
<b>Summary estimate (95% CI)*</b>					<b>0.46 (0.26–0.80)</b>	<b>97.7% (96.1–98.7%)</b>
<b>Retrospective studies</b>						
Schneider et al. (2011)	100	6.2	6	620	0.97	95.3
Krennmair et al. (2010)	112	5.7	4	642	0.62	96.9
Urdaneta et al. (2010)	326	5.9	6	1921	0.31	98.5
Zafropoulos et al. (2010)	252	4.8	11	1205	0.91	95.5
Hälg et al. (2008)	22	5	1	111	0.9	95.6
Pikner et al. (2008)	45	5	1	225	0.44	97.8
De Boever et al. (2006)	80	5.2	0	417	0	100.0
Wagenberg & Froum (2006)	401	5.9	18	2266	0.79	96.1
Elkhoury et al. (2005)	39	5	0	195	0	100.0
Levin et al. (2005)	30	5.1	2	153	1.31	93.7
Bernard et al. (2004)	32	5	0	158	0	100.0
Vigolo & Givani (2000)	52	5	3	245	1.22	94.1
<b>Total</b>	<b>1491</b>	<b>5.3</b>	<b>52</b>	<b>8158</b>		
<b>Summary estimate (95% CI)*</b>					<b>0.78 (0.28–2.20)</b>	<b>96.2% (89.6–98.6%)</b>
<b>Total</b>	<b>2803</b>	<b>5.2</b>	<b>83</b>	<b>14715</b>		
<b>Overall summary estimate (95% CI)*</b>					<b>0.56 (0.42–0.76)</b>	<b>97.2% (96.3–97.9%)</b>

\*Based on standard Poisson regression, test for heterogeneity  $P = 0.141$ 

implants); Forty-nine implants were lost after loading (1.5% of all placed implants). In one study, only the number of implants lost in function ( $n = 2$ ) was reported (Schmidlin et al. 2010), whereas two studies did not specify the time-point of implant failure (Krieger et al. 2009; Zafropoulos et al. 2010). For failures after loading, the estimated annual failure rates were 0.29 (95% CI: 0.17–0.47; 36 studies) over 5 years and 0.35 (95% CI: 0.15–0.83; 10 studies) over 10 years.

The study-specific 5-year survival proportion varied between 90.5% and 100% (Table 3) with an estimated failure rate per 100 implant years between 0 and 2 (Fig. 2). Similar calculations for the 10-year survival proportion ranged between 85.5% and 100% (Table 4), whereas the estimated failure rate

per 100 implant years was between 0 and 1.56 (Fig. 3). Based on the meta-analysis, this estimated failure rate per 100 implant years resulted in 0.56 (95% CI: 0.42–0.76; all 36 studies; Fig. 2), 0.46 (95% CI: 0.26–0.80; 23 prospective studies), and 0.78 (95% CI: 0.28–2.20; 13 retrospective studies) over 5 years (Table 3), and in 0.49 (95% CI: 0.28–0.85; all 10 studies; Fig. 3), 0.52 (95% CI: 0.25–1.08; 4 prospective studies), and 0.48 (95% CI: 0.22–1.03; 6 retrospective studies) over 10 years (Table 4). The respective implant survival rates for implants supporting SCs at 5 years amounted to 97.2% (95% CI: 96.3–97.9%; all 36 studies), 97.7% (95% CI: 96.1–98.7%; 4 prospective studies), and 96.2% (95% CI: 89.6–98.6%; 6 retrospective studies) (Table 3), and at 10 years to 95.2% (95% CI: 91.8–97.2%; all

36 studies), 94.9% (95% CI: 89.7–97.5%; 4 prospective studies), and 95.3% (95% CI: 90.2–97.8%; 6 retrospective studies) (Table 4).

### SC survival

The survival of SC was defined as SCs remaining *in situ* with or without modification during the observation period. Twenty studies provided data with a mean follow-up of 5 years and a total number of 1385 SCs (Table 5). Of 1385 SCs, 53 crowns were lost, resulting in a study-specific 5-year survival rate between 89.6% and 100% (Table 5). Twenty-eight SCs were lost because the implants were lost, whereas in 25 SCs only the reconstructions failed. The failure rate per 100 SC years ranged between 0 and 2.19 (Table 5). The meta-analysis demonstrated an

**Table 4. Annual failure rates and 10-year survival of implants**

Study	Total number of implants	Mean follow-up time	Number of failure	Total implant exposure time	Estimated failure rate (per 100 implant years)	Estimated survival rate after 10 years (in percent)
<b>Prospective Studies</b>						
MacDonald et al. (2009)	20	7.7	1	154	0.65	93.7
Gotfredsen (2009)	20	10	0	200	0	100.0
Brägger et al. (2005)	69	10	5	672	0.74	97.6
Thilander et al. (1999)	15	8	0	120	0	100.0
<b>Total</b>	<b>124</b>	<b>8.9</b>	<b>6</b>	<b>1146</b>		
<b>Summary estimate (95% CI)*</b>					<b>0.52 (0.25–1.08)</b>	<b>94.9% (89.7–97.5%)</b>
<b>Retrospective Studies</b>						
Jung et al. (2012a)	20	12.5	1	250	0.4	96.1
Bonde et al. (2010)	55	9.4	3	515	0.58	94.3
Matarasso et al. (2010)	80	9.7	6	773	0.78	92.5
Krieger et al. (2009)	24	8	3	192	1.56	85.5
Jemt (2009)	41	10	0	410	0	100.0
Jemt (2008)	47	12.3	0	576	0	100.0
<b>Total</b>	<b>267</b>	<b>10.3</b>	<b>13</b>	<b>2716</b>		
<b>Summary estimate (95% CI)*</b>					<b>0.48 (0.22–1.03)</b>	<b>95.3% (90.2–97.8%)</b>
<b>Total</b>	<b>391</b>	<b>9.9</b>	<b>19</b>	<b>3862</b>		
<b>Overall summary estimate (95% CI) *</b>					<b>0.49 (0.28–0.85)</b>	<b>95.2% (91.8–97.2%)</b>

\*Based on standard Poisson regression, test for heterogeneity  $P = 0.152$

annual failure rate of 0.75 (95% CI: 0.48 – 1.18; all 20 studies; Fig. 4), 0.76 (95% CI: 0.38–1.54; 14 prospective studies), and 0.68 (95% CI: 0.41–1.10; 6 retrospective studies) (Table 5). This translated into a survival rate for implant-supported SCs of 96.3% (95% CI: 94.2–97.6%; all 20 studies), 96.5% (95% CI: 92.6–98.1%; 14 prospective studies), and of 96.7% (95% CI: 94.6–97.7%; 6 retrospective studies) after 5 years (Table 5). Similar calculations were performed for studies with a mean observation period of 10 years and included seven studies and 268 SCs (Table 6).

Twenty-eight failures were reported (8 in combination with implant failure; 20 failure of the reconstruction only). The failure rate per 100 SC years ranged from 0.58 to 2.19. The meta-analysis revealed an annual failure rate of 1.12 (95% CI: 0.66 – 1.89; all 7 studies; Fig. 5), 1.07 (95% CI: 0.97–1.19; 3 prospective studies), and 1.14 (95% CI: 0.48–2.73; 4 retrospective studies) (Table 5). The calculated survival rate for implant-supported SCs was 89.4% (95% CI: 82.8–93.6%; all seven studies), 89.8% (95% CI: 88.8–90.8%; three prospective studies), and of 89.2% (95%

CI: 76.1–95.3%; four retrospective studies) after 10 years (Table 6).

In addition, multivariate Poisson regression was applied to account for the influence of the type of fixation on the survival rate of SCs. The calculated survival rate of cemented SCs (15 studies, 872 crowns) was 95.6% (95% CI: 93.0–97.2%) and 95.0% (95% CI: 92.1–96.9%) for screw-retained SCs (5 studies, 545 crowns). This difference was not statistically significant ( $P > 0.05$ ).

To take into account the reconstruction materials, studies were also divided into groups with metal-ceramic crowns (17 studies, 799 SCs) and all-ceramic crowns (2 studies, 117 SCs) (Andersson et al. 1998a; Bonde et al. 2010). The stratified summary estimated of the survival proportion after 5 years of loading amounted to 95.8% (95% CI: 93.1–97.5%) for metal-ceramic crowns and 95.8% (95% CI: 90.7–98.1%) for all-ceramic crowns. The annual failure rates of 0.85 (95% CI: 0.51–1.42) for metal-ceramic crowns and 0.86 (95% CI: 0.38–1.95) for all-ceramic crowns did not reveal statistical significance ( $P > 0.05$ ) based on standard Poisson regression.

**Biological outcomes**

Biological complications were reported in 15 studies and included various descriptions of any kind of soft tissue complications: signs of inflammation, mucosal inflammation, mucositis, bleeding, suppuration, and soft tissue dehiscences. The meta-analysis revealed an estimated rate of various types of soft tissue complications (per 100 implant years) of 1.47 (95% CI: 0.90–2.39). This resulted in a 5-year cumulative soft tissue complication rate of 7.1% (95% CI: 4.4–11.3%) (Table 7).




Fig. 2. Annual failure rates (per 100 years) of implants after 5 years.




Fig. 3. Annual failure rates (per 100 years) of implants after 10 years.

In 17 studies, a radiographic analysis was performed to evaluate the peri-implant bone levels. Based on the meta-analysis, the cumulative 5-year complication rate (implants

with bone loss >2 mm) was 5.2% (95% CI: 3.1–8.6%). The estimated rate of bone loss >2 mm per 100 implant years amounted to 1.06 (95% CI: 0.62–1.79) (Table 7).

Multivariate Poisson regression was used to account for the influence of the type of fixation of the reconstruction (cemented, screw-retained) on marginal bone loss >2 mm. The 5-year complication rate (implants with bone loss >2 mm) was slightly higher for cemented reconstructions (2.8%; 95% CI: 2.1–3.7%) than that for screw-retained reconstructions (1.1%; 95% CI: 0.2–7.1%). However, the annual complication rate of 0.56 (95% CI: 0.42–0.76) for cemented SCs and 0.22 (95% CI: 0.03–1.46) for screw-retained SCs did not reveal a statistically significant influence of the crown design ( $P > 0.05$ ).

**Aesthetic outcomes**

A variety of studies reported on aesthetic outcomes. Outcome measures were evaluated by dental professionals or by patients and included the use of a questionnaire to rate the appearance of the crown or an index system to rate the interdental papilla height (Schropp & Isidor 2008; Gotfredsen 2009; MacDonald et al. 2009; Krennmair et al. 2010). Twelve studies reported on aesthetic complications with crowns having a semi-optimal or even an unacceptable aesthetic appearance due to soft tissue recessions, an unfavorable color, and visible crown margins. The cumulative

Table 5. Annual failure rates and 5-year survival of implant-supported SCs

Study	Total number of single crowns	Mean follow-up time	Number of failure	Total crown exposure time	Estimated failure rate (per 100 crown years)	Estimated survival rate after 5 years (in percent)
<b>Prospective Studies</b>						
Jung et al. (2012b)	29	4.7	0	137	0	100.0
Vigolo & Givani (2009)	182	5	0	910	0	100.0
Degidi et al. (2008)	45	5	0	225	0	100.0
Schropp & Isidor (2008)	42	4.7	2	200	1	95.1
Kreissl et al. (2007)	46	5	1	230	0.43	97.8
Wennström et al. (2005)	44	5	1	208	0.48	97.6
Gotfredsen (2004)	20	5	1	98	1.02	95.0
Haas et al. (2002)	75	5.5	4	382	1.05	94.9
Mericske-Stern et al. (2001)	26	6.5	2	169	1.18	94.3
Palmer et al. (2000)	15	5	1	66	1.52	92.7
Polizzi et al. (1999)	30	5.3	2	154	1.3	93.7
Andersson et al. (1998a)	38	5	1	179	0.56	97.2
Andersson et al. (1998b)	65	5	4	295	1.36	93.4
Scheller et al. (1998)	97	5	9	411	2.19	89.6
<b>Total</b>	<b>754</b>	<b>5.1</b>	<b>28</b>	<b>3664</b>		
<b>Summary estimate (95% CI)*</b>					<b>0.76 (0.38–1.54)</b>	<b>96.3% (92.6–98.1%)</b>
<b>Retrospective Studies</b>						
Schneider et al. (2011)	100	6.2	6	620	0.97	95.3
Krennmair et al. (2010)	112	5.7	0	642	0	100.0
Schmidlin et al. (2010)	39	6.2	2	243	0.82	96.0
Urdaneta et al. (2010)	326	5.9	16	1921	0.83	95.9
Hälg et al. (2008)	22	5	1	111	0.9	95.6
Bernard et al. (2004)	32	5	0	158	0	100.0
<b>Total</b>	<b>631</b>	<b>5.7</b>	<b>25</b>	<b>3695</b>		
<b>Summary estimate (95% CI)*</b>					<b>0.68 (0.41–1.10)</b>	<b>96.7% (94.6–97.7%)</b>
<b>Total</b>	<b>1385</b>	<b>5.3</b>	<b>53</b>	<b>7359</b>		
<b>Overall summary estimate (95% CI)*</b>					<b>0.75 (0.48–1.18)</b>	<b>96.3% (94.2–97.6%)</b>

\*Based on standard Poisson regression, test for heterogeneity  $P = 0.024$






Fig. 4. Annual failure rates (per 100 years) of SCs after 5 years.




Fig. 5. Annual failure rates (per 100 years) of SCs after 10 years.

5-year aesthetic complication rate was 7.1% (95% CI: 3.6–13.6%) (Table 7).

**Technical outcomes**

A variety of technical complications were reported in 36 studies. The most common technical complication was abutment- or screw-loosening, reaching a cumulative incidence of 8.8% (95% CI: 5.1–15.0%) after

5 years (Table 8). Although no such complications were reported in six studies, one study was a clear outlier with an estimated rate of 18.03 (Henry et al. 1996). In that study, SCs were mounted on Brånemark implants using gold screws.

The second most common technical complication was loss of retention (fracture of the luting cement), reported in 13 studies and

reaching a cumulative incidence of 4.1% (95% CI: 2.2–7.5%) after 5 years (Table 8).

The third most common technical complication was reported for fracture of the veneering material (acrylic or ceramic chippings). The cumulative complication rate amounted to 3.5% (95% CI: 2.4–5.2%) after 5 years. These incidences include minor (chippings that can be polished) and major (repair necessary) fractures of the veneering material. No statistically significant differences with respect to the incidence of veneer fractures were observed between porcelain fused to metal crowns and all-ceramic crowns ( $P > 0.05$ ). Fracture of the framework material was reported in 16 studies, but it only happened in 6 studies, resulting in a cumulative complication rate of 3.5% (95% CI: 2.4–4.1%) after 5 years. The incidence of framework fractures between porcelain fused to metal crowns and all-ceramic crowns was not statistically significantly different ( $P > 0.05$ ).

Additional technical complications were rarely observed. The cumulative 5-year complication rates amounted to 0.18% (95% CI: 0.03–0.4%) for implant fractures and to 0.18% (95% CI: 0.03–0.4%) for abutment or screw fractures. Loss of the access hole restoration was never observed and only reported by three studies.

**Discussion**

This systematic review addressed the survival and complication rates of implant-supported SCs based on clinical studies with a mean observation period of at least 5 years. The outcomes of the meta-analysis demonstrated both high implant survival rates for single tooth implants and high survival rates of the respective SCs after 5 and 10 years. It must be noted, however, that the most common complications reached 8.8% (technical), 7.1% (biological), and 7.1% (aesthetic) over 5 years.

**Implant survival**

The calculated implant survival based on 46 included studies with a mean observation period of 5 years amounted 97.2% (95% CI: 96.3–97.9%) after 5 years and 95.2% (95% CI: 91.8–97.2%) after 10 years. The implant survival rate at 5 years was based on 2803 implants and even slightly higher than that in the previous systematic review (96.8%) with a lower number of implants (1558) (Jung et al. 2008a). Based on 10 studies, the estimated implant survival rate after 10 years

**Table 6. Annual failure rates and 10-year survival of implant-supported SCs**

Study	Total number of single crowns	Mean follow-up time	Number of failure	Total crown exposure time	Estimated failure rate (per 100 crown years)	Estimated survival rate after 5 years (in percent)
<b>Prospective Studies</b>						
Gotfredsen (2009)	20	10	2	190	1.05	90.0
Brägger et al. (2005)	69	10	7	623	1.12	89.4
Thilander et al. (1999)	15	8	1	120	0.83	92.0
<b>Total</b>	<b>104</b>	<b>9.3</b>	<b>10</b>	<b>933</b>		
<b>Summary estimate (95% CI)*</b>					<b>1.07 (0.97–1.19)</b>	<b>89.8% (88.8–90.8%)</b>
<b>Retrospective Studies</b>						
Bonde et al. (2010)	52	9.4	3	515	0.58	94.3
Krieger et al. (2009)	24	8	4	192	2.08	81.2
Jemt (2009)	41	10	1	410	0.24	97.6
Jemt (2008)	47	12.3	10	457	2.19	80.3
<b>Total</b>	<b>164</b>	<b>9.9</b>	<b>18</b>	<b>1574</b>		
<b>Summary estimate (95% CI)*</b>					<b>1.14 (0.48–2.73)</b>	<b>89.2% (76.1–95.3%)</b>
<b>Total</b>	<b>268</b>	<b>9.4</b>	<b>28</b>	<b>2507</b>		
<b>Overall summary estimate (95% CI)*</b>					<b>1.12 (0.66–1.89)</b>	<b>89.4% (82.8–93.6%)</b>

\*Based on standard Poisson regression, test for heterogeneity  $P = 0.105$

could be calculated, and revealed an even lower estimated annual failure rate (per 100 implant years) of 0.49 after 10 years compared to 0.56 after 5 years. It is not uncommon to observe higher implant failure rates in shorter termed studies, because in this review roughly half of all implant failures were early failures before loading. With a respective 10-year implant survival rate of 95.2% this treatment modality can be considered as safe and predictable. The lowest implant survival rate of all included studies was 85.5% after 10 years (Krieger et al. 2009). This particular retrospective study exclusively reported on patients with birth defects affecting the formation of teeth. They concluded that especially in cases with cleft lip, alveolus, and palate (CLAP), in which anatomical conditions render implant placement difficult and in which teeth adjacent to the cleft require aesthetic corrections, the conventional FDP on teeth might be the treatment of choice.

#### SC survival

In this systematic review, the survival rate for implant-supported SCs was 96.3% after 5 years of loading. This value is slightly higher compared to the results of the previous systematic review reporting a survival

rate of 94.5% after 5 years for implant-supported SCs (Jung et al. 2008a). Based on this comparison and on Table 5, a trend can be recognized with newer studies reporting higher survival rates for implant-supported SCs. Consequently, the highest failure rate after 5 years (10.4%) was reported by the oldest included study within this systematic review (Scheller et al. 1998). This trend might be explained by the fact that newer studies included implant systems with improved implant types and designs as well as enhanced prosthetic components, which may allow reducing the incidence of failures.

After 10 years, this meta-analysis reveals a survival rate for implant-supported SCs of 89.4% derived from seven studies including 268 implant-supported SCs. This is an important number when it comes to the decision-making process between the different treatment modalities for a single tooth gap. Hence, the 10-year outcome for the implant-supported SCs must be compared to the outcomes of conventional and cantilever FDPs. The meta-analysis of conventional FDPs indicated an estimated survival rate of 89.1% (95% CI: 81.0–93.8%) after 10 years (Tan et al. 2004). The estimated survival rate of cantilever FDPs was 81.8% (95% CI: 78.2–84.9%) after 10 years (Pjetursson et al. 2004). This compar-

ison of the survival rates after 10 years demonstrates that the calculated numbers for the implant-supported SCs are very similar to the ones from the conventional FDPs and more favorable compared to cantilever FDPs.

#### Biological outcomes

Biological complications have been reported in the dental literature very inconsistently and without any standardization and classification. This results in a large variety of clinical reports ranging from signs of inflammation, mucosa inflammation, mucositis, bleeding, and suppuration to soft tissue dehiscences. Summarizing all these complications independent of their severity, the cumulative soft tissue complication rate was 7.1% after 5 years. Compared to the previous systematic review demonstrating a soft tissue complication rate of 9.7% after 5 years, there is also a trend to less soft tissue complications when more and especially newer studies are analyzed (Jung et al. 2008a).

Looking at the cumulative 5-year complication rates of implants with bone loss exceeding 2 mm can identify the same trend. This review revealed a complication rate of 5.2% after 5 years, whereas in the former review a complication rate of implants having bone loss >2 mm of 6.3% was calculated (Jung et al. 2008a).

The type of fixation of the reconstruction (cemented, screw-retained) did not have any significant influence on the estimated rate of biological complications ( $P > 0.05$ ).

#### Aesthetic outcomes

The aesthetic outcome has certainly been not only considered as the major focus from a patient's perspective but also from the clinician's side. Currently available indices to rate the aesthetic outcomes of SCs include measurements of the papilla height and questionnaires for patients and lay persons (Jemt 1999; Schropp & Isidor 2008; Gotfredsen 2009; MacDonald et al. 2009; Krennmair et al. 2010). In this systematic review, a variety of the included publications reported on aesthetic complications (e.g., dehiscences of the soft tissue with exposure of the crown margin, suboptimal color of the prosthetic reconstruction) and on general aesthetic outcomes (e.g., papilla height measurements, questionnaires). However, due to a lack of standardized parameters and indices to evaluate the aesthetic appearance, a large heterogeneity exists between the different studies. This may limit the scientific value of the calculated cumulative 5-year aesthetic complication rate of 7.1%, because this is based on various

**Table 7. Biological and aesthetic complications**

Study	Total number of implants	Total implant exposure time	Estimated rate of bone loss >2 mm (per 100 implant years)	Estimated rate of soft tissue complications (per 100 implant years)	Estimated rate of aesthetic complications (per 100 crown years)
Jung et al. (2012a)	20	250	1.2	NR	1.29
Schneider et al. (2011)	100	620	1.77	0	NR
Bonde et al. (2010)	55	515	NR	1.36	NR
Matarasso et al. (2010)	80	773	3.49	NR	NR
Schmidlin et al. (2010)	39	243	NR	2.06	0
MacDonald et al. (2009)	20	154	0.65	NR	NR
Gottfredsen (2009)	20	200	0.5	1	NR
Hälg et al. (2008)	22	111	0	NR	NR
Jemt (2009)	41	410	0.49	0.98	0.49
Jemt (2008)	47	576	0	2.26	1.97
Schropp & Isidor (2008)	45	210	0.95	1.43	1
Bornstein et al. (2005)	39	190	0	0	NR
Elkhoury et al. (2005)	39	195	3.08	NR	NR
De Boever & deBoever (2005)	10	50	2	NR	NR
Wennström et al. (2005)	45	208	0.96	NR	NR
Levin et al. (2005)	51	195	NR	NR	3.59
Jemt & Lekholm (2005)	10	48	0	NR	NR
Brägger et al. (2005)	69	672	NR	1.93	NR
Bernard et al. (2004)	32	158	0	NR	0
Gottfredsen (2004)	20	100	NR	NA	NR
Haas et al. (2002)	76	382	NR	NR	0.52
Gibbard & Zarb (2002)	49	287	NR	1.05	1.05
Mericske-Stern et al. (2001)	26	169	0.59	NR	NR
Palmer et al. (2000)	15	70	NR	0	NR
Andersson et al. (1998a)	38	182	1.1	1.1	0.56
Andersson et al. (1998b)	65	305	NR	0.33	0
Scheller et al. (1998)	99	411	NR	1.22	NR
Henry et al. (1996)	107	477	NR	6.08	6.71
<b>Summary estimate event rates (95% CI)</b>			<b>1.06* (0.62–1.79)</b>	<b>1.47* (0.90–2.39)</b>	<b>1.47* (0.74–2.92)</b>
<b>Cumulative 5 year complication rates (95% CI)</b>			<b>5.2%* (3.1–8.6%)</b>	<b>7.1%* (4.4–11.3%)</b>	<b>7.1%* (3.6–13.6%)</b>

NR, not reported; NA, not analyzed;  
\*Based on random-effects Poisson regression.

measurements and parameters. A scientific consensus on an accepted and reproducible method to evaluate the aesthetic outcome of SCs on the soft tissue level and on the level of the crown itself would therefore be needed.

#### Technical outcomes

In agreement with previous systematic reviews, this study also revealed that abutment- or screw-loosening are the most common technical complications (Berghlundh et al. 2002; Jung et al. 2008a; Sailer et al. 2009). For implant-supported SCs the incidence of abutment or screw-loosening was 8.8% after 5 years. However, it must be emphasized that two studies using an old gold-screw design were mainly responsible for the high number of screw-loosening (Henry et al. 1996; Jemt 2008).

When it comes to the comparison of all-ceramic vs. porcelain-fused to metal (PFM) crowns, the overall survival rate, the fracture rate of the veneering ceramic, and the incidence of framework fractures are of primary interest. The type of the reconstruction did not influence the survival rate of SCs based on standard Poisson regression. This is in con-

trast to the previously published systematic review (Jung et al. 2008a), but in agreement with a more recently published systematic review focusing specifically on the comparison between metal-ceramic and all-ceramic reconstructions (Sailer et al. 2009). In the latter, no statistically significant differences were found between metal-ceramic and all-ceramic crowns based on the calculated estimated 5-year cumulative survival rate (Sailer et al. 2009). The overall cumulative fracture rate of the veneering material amounted to 3.5% after 5 years with no statistically significant differences between all-ceramic and PFM crowns. This was also true for the incidence of framework fractures (3.5% after 5 years) with no significant difference between all-ceramic and PFM crowns. This is confirmed by a recent systematic review comparing the performance of all-ceramic and metal abutments and the corresponding reconstructions (Sailer et al. 2009). They provided no statistically significant differences for technical complications of ceramic and metal abutments after at least 3 years. However, it was emphasized that the information for ceramic abutments was limited in the number of studies and

abutments analyzed as well as the accrued follow-up time.

## Conclusion

The outcomes of the meta-analysis demonstrated both, high implant survival rates for single tooth implants and the respective single crowns after 5 and 10 years. Despite varying rates of technical, biological, and aesthetic complications that need to be expected, this treatment modality for the restoration of a single tooth gap can be considered as a safe and predictable therapeutic option.

#### Clinical recommendations

Considering high implant and SC survival rates observed in prospective and retrospective studies with a mean follow-up of 5 and 10 years, this treatment modality can be recommended for single tooth gaps. Clinicians must be aware that complications may occur to various extents. Most notably, abutment and screw-loosening were reported with the highest technical complications. Although the dental literature reports soft tissue and

**Table 8. Technical complications**

Study	Total number of implants	Estimated rate of implant fracture (per 100 implant years)	Total number of crowns	Estimated rate of abutment or screw fracture (per 100 crown years)	Estimated rate of loose abutments or screws (per 100 crown years)	Estimated rate of loss of retention (per 100 crown years)	Estimated rate of ceramic chipping (per 100 crown years)	Estimated rate of framework fracture (per 100 crown years)	Estimated rate of loss of access hole restoration (per 100 crown years)
Schneider et al. (2011)	100	0	100	0	1.29	0.81	0.65	0	0
Bonde et al. (2010)	55	0	52	0	0.58	NR	0.58	0.19	NR
Krennmair et al. (2010)	112	0	112	0	0.78	1.71	0.78	0	NR
Schmidlin et al. (2010)	39	0	39	0	0.82	0.41	0.82	0	0
Urdaneta et al. (2010)	326	0	326	0.16	0.94	NR	0.94	NR	NR
MacDonald et al. (2009)	20	0	20	0	1.3	NR	0	0	0
Vigolo & Givani (2009)	182	0	182	0	0	0	0	1.39	NR
Gotfredsen (2009)	20	0	20	0	1.05	1.05	1.58	0	NR
Hälg et al. (2008)	22	0.9	22	0	0	0	0	0	NR
Jemt (2009)	41	0	41	0	1.22	NR	NR	NR	NR
Jemt (2008)	47	0	47	NR	6.13	NR	NR	NR	NR
Schropp & Isidor (2008)	45	0	42	0	0	1.5	0	0	NR
Kreissl et al. (2007)	46	0	46	0.87	2.61	NR	0.87	0	NR
Romeo et al. (2006)	58	0	58	NR	NR	0.36	1.39	NR	NR
Wagenberg & Froum (2006)	401	0	383	NR	NR	NR	NR	NR	NR
Bornstein et al. (2005)	39	0	39	NR	NR	NR	NR	NR	NR
Elkhoury et al. (2005)	39	0	39	NR	NR	NR	NR	NR	NR
De Boever & de Boever (2005)	10	0	10	NR	NR	NR	NR	NR	NR
Wennström et al. (2005)	45	0	44	0	1.44	NR	NR	0	NR
Jemt & Lekholm (2005)	10	0	10	NR	NR	NR	NR	NR	NR
Brägger et al. (2005)	69	NR	69	0	0.48	0	0.48	0	NR
Taylor et al. (2004)	39	0	38	NR	NR	NR	NR	NR	NR
Bernard et al. (2004)	32	0	32	NR	NR	NR	NR	NR	NR
Romeo et al. (2004)	123	0	121	0	0	0.56	0.28	NR	NR
Bianchi & Sanfilippo (2004)	94	0	94	NR	NR	NR	NR	NR	NR
Gotfredsen (2004)	20	0	20	NR	NA	NA	NA	NA	NR
Haas et al. (2002)	76	0.26	77	NR	3.14	NR	NR	NR	NR
Gibbard & Zarb (2002)	49	0	48	NR	1.39	NR	NR	NR	NR
Palmer et al. (2000)	15	0	15	0	0	1.52	NR	1.52	NR
Vigolo & Givani (2000)	52	0	52	0	0.41	2.86	NR	NR	NR
Thilander et al. (1999)	15	0	15	NR	NR	NR	NR	NR	NR
Polizzi et al. (1999)	30	0.63	30	0.65	0	NR	0	NR	NR
Andersson et al. (1998a)	38	0	38	NR	NR	NR	NR	0.56	NR
Andersson et al. (1998b)	65	0	65	0	0.34	NR	0.34	0.68	NR
Scheller et al. (1998)	99	NR	97	NR	0.97	0.73	1.7	1.7	NR
Henry et al. (1996)	107	0	106	0.21	18.03	NR	1.89	NR	NR
<b>Summary estimate event rates (95% CI)</b>		<b>0.03* (0.007–0.19)</b>		<b>0.08* (0.027–0.23)</b>	<b>1.84* (1.04–3.25)</b>	<b>0.84* (0.45–1.56)</b>	<b>0.72* (0.48–1.08)</b>	<b>0.26* (0.08–0.84)</b>	<b>0 (0–0.36)</b>
<b>Cumulative 5 year complication rates (95% CI)</b>		<b>0.18%* (0.03–0.4%)</b>		<b>0.4%* (0.14–1.1%)</b>	<b>8.8%* (5.1–15.0%)</b>	<b>4.1%* (2.2–7.5%)</b>	<b>3.5%* (2.4–5.2%)</b>	<b>1.3%* (0.4–4.1%)</b>	<b>0%</b>

NR, not reported; NA, not analyzed;  
\*Based on random-effects Poisson regression.

aesthetic complications very inconsistently and without any standardization and classification, these complications have to be considered and strengthen the need for a well-established maintenance program.

### Research recommendations

The outcomes of this systematic review are based on a large variety of studies with differing levels of evidence. Although basic biological parameters (e.g., marginal bone levels)

were frequently reported in the studies, technical outcome measures were inconsistently analyzed. In addition, many of the long-term studies include reconstruction materials that are no longer in use. It is therefore of great interest to perform prospective long-term studies evaluating current implant types, design, and prosthetic components and to assure that standardized technical, biological, and aesthetic outcome measures are used.

### Acknowledgements

The authors are grateful to Gisela Müller, Study Monitor, Clinic of Fixed and Removable Prosthodontics and Dental Material Science, University of Zurich, for her help in the preparation of the manuscript.

### References

- Andersson, B., Odman, P., Lindvall, A.M. & Brånemark, P.I. (1998b) Cemented single crowns on osseointegrated implants after 5 years: results from a prospective study on CeraOne. *The International Journal of Prosthodontics* **11**: 212–218.
- Berglundh, T., Persson, L. & Klinge, B. (2002) A systematic review of the incidence of biological and technical complications in implant dentistry reported in prospective longitudinal studies of at least 5 years. *Journal of Clinical Periodontology* **29**(Suppl 3): 197–212; discussion 232–193.
- Bonde, M.J., Stokholm, R., Isidor, F. & Schou, S. (2010) Outcome of implant-supported single-tooth replacements performed by dental students. A 10-year clinical and radiographic retrospective study. *European Journal of Oral Implantology* **3**: 37–46.
- Egger, M., Smith, G.D. & Sterne, J.A. (2001) Uses and abuses of meta-analysis. *Clin Med* **1**: 478–484.
- Gottfredsen, K. (2009) A 10-Year Prospective Study of Single Tooth Implants Placed in the Anterior Maxilla. *Clinical Implant Dentistry and Related Research*. Epub August 14, 2009.
- Henry, P.J., Laney, W.R., Jemt, T., Harris, D., Krogh, P.H., Polizzi, G., Zarb, G.A. & Herrmann, I. (1996) Osseointegrated implants for single-tooth replacement: a prospective 5-year multicenter study. *The International Journal of Oral & Maxillofacial Implants* **11**: 450–455.
- Jemt, T. (1999) Restoring the gingival contour by means of provisional resin crowns after single-implant treatment. *International Journal of Periodontics and Restorative Dentistry* **19**: 20–29.
- Jemt, T. (2008) Single implants in the anterior maxilla after 15 years of follow-up: comparison with central implants in the edentulous maxilla. *The International Journal of Prosthodontics* **21**: 400–408.
- Jung, R.E., Pjetursson, B.E., Glauser, R., Zembic, A., Zwahlen, M. & Lang, N.P. (2008a) A systematic review of the 5-year survival and complication rates of implant-supported single crowns. *Clinical Oral Implants Research* **19**: 119–130.
- Kirkwood, B.R. & Sterne, J.A.C. (2003a) Survival analysis: displaying and comparing survival patterns. *Essential Medical Statistics*, pp. 272–286.
- Kirkwood, B.R. & Sterne, J.A.C. (2003b) Poisson regression. *Essential Medical Statistics*. pp. 249–262. Oxford: Blackwell Science Ltd.
- Krennmair, G., Seemann, R., Schmidinger, S., Ewers, R. & Piehslinger, E. (2010) Clinical outcome of root-shaped dental implants of various diameters: 5-year results. *The International Journal of Oral & Maxillofacial Implants* **25**: 357–366.
- Krieger, O., Matulienė, G., Husler, J., Salvi, G.E., Pjetursson, B. & Bragger, U. (2009) Failures and complications in patients with birth defects restored with fixed dental prostheses and single crowns on teeth and/or implants. *Clinical Oral Implants Research* **20**: 809–816.
- Levin, L., Pathael, S., Dolev, E. & Schwartz-Arad, D. (2005) Aesthetic versus surgical success of single tooth implants: 1- to 9-year follow-up. *Practical procedures & aesthetic dentistry: PPAD* **17**: 533–538; quiz 540, 566.
- MacDonald, K., Pharoah, M., Todescan, R. & Deporter, D. (2009) Use of sintered porous-surfaced dental implants to restore single teeth in the maxilla: a 7- to 9-year follow-up. *The International Journal of Periodontics & Restorative Dentistry* **29**: 191–199.
- Pjetursson, B.E., Tan, K., Lang, N.P., Bragger, U., Egger, M. & Zwahlen, M. (2004) A systematic review of the survival and complication rates of fixed partial dentures (FPDs) after an observation period of at least 5 years. *Clinical Oral Implants Research* **15**: 625–642.
- Sailer, I., Philipp, A., Zembic, A., Pjetursson, B.E., Hammerle, C.H. & Zwahlen, M. (2009) A systematic review of the performance of ceramic and metal implant abutments supporting fixed implant reconstructions. *Clinical Oral Implants Research* **20**(Suppl 4): 4–31.
- Scheller, H., Urgell, J.P., Kultje, C., Klineberg, I., Goldberg, P.V., Stevenson-Moore, P., Alonso, J. M., Schaller, M., Corria, R.M., Engquist, B., Toreskog, S., Kastenbaum, F. & Smith, C.R. (1998) A 5-year multicenter study on implant-supported single crown restorations. *The International Journal of Oral & Maxillofacial Implants* **13**: 212–218.
- Schmidlin, K., Schnell, N., Steiner, S., Salvi, G.E., Pjetursson, B., Matulienė, G., Zwahlen, M., Bragger, U. & Lang, N.P. (2010) Complication and failure rates in patients treated for chronic periodontitis and restored with single crowns on teeth and/or implants. *Clinical Oral Implants Research* **21**: 550–557.
- Schropp, L. & Isidor, F. (2008) Clinical outcome and patient satisfaction following full-flap elevation for early and delayed placement of single-tooth implants: a 5-year randomized study. *The International Journal of Oral & Maxillofacial Implants* **23**: 733–743.
- Tan, K., Pjetursson, B.E., Lang, N.P. & Chan, E.S. (2004) A systematic review of the survival and complication rates of fixed partial dentures (FPDs) after an observation period of at least 5 years. *Clinical Oral Implants Research* **15**: 654–666.
- Zafropoulos, G.G., Deli, G., Bartee, B.K. & Hoffmann, O. (2010) Single-tooth implant placement and loading in fresh and regenerated extraction sockets. Five-year results: a case series using two different implant designs. *Journal of Periodontology* **81**: 604–615.
- Al-Amleh, B., Lyons, K. & Swain, M. (2010) Clinical trials in zirconia: a systematic review. *Journal of Oral Rehabilitation* **37**: 641–652.
- Andreietelli, M., Wenz, H.J. & Kohal, R.J. (2009) Are ceramic implants a viable alternative to titanium implants? A systematic literature review. *Clinical Oral Implants Research* **20**(Suppl 4): 32–47.
- Atieh, M.A., Atieh, A.H., Payne, A.G. & Duncan, W.J. (2009a) Immediate loading with single implant crowns: a systematic review and meta-analysis. *International Journal of Prosthodontics* **22**: 378–387.
- Atieh, M.A., Payne, A.G., Duncan, W.J. & Cullinan, M.P. (2009b) Immediate restoration/loading of immediately placed single implants: is it an effective bimodal approach? *Clinical Oral Implants Research* **20**: 645–659.
- Atieh, M.A., Ibrahim, H.M. & Atieh, A.H. (2010a) Platform switching for marginal bone preservation around dental implants: a systematic review and meta-analysis. *Journal of Periodontology* **81**: 1350–1366.
- Atieh, M.A., Payne, A.G., Duncan, W.J., de Silva, R.K. & Cullinan, M.P. (2010b) Immediate placement or immediate restoration/loading of single implants for molar tooth replacement: a systematic review and meta-analysis. *International Journal of Oral and Maxillofacial Implants* **25**: 401–415.

- Blanes, R.J. (2009) To what extent does the crown-implant ratio affect the survival and complications of implant-supported reconstructions? A systematic review. *Clinical Oral Implants Research* **20**(Suppl 4): 67–72.
- Brodala, N. (2009) Flapless surgery and its effect on dental implant outcomes. *International Journal of Oral and Maxillofacial Implants* **24**(Suppl): 118–125.
- Chen, S.T. & Buser, D. (2009) Clinical and esthetic outcomes of implants placed in postextraction sites. *International Journal of Oral and Maxillofacial Implants* **24**(Suppl): 186–217.
- Cordaro, L., Torsello, F. & Rocuzzo, M. (2009a) Implant loading protocols for the partially edentulous posterior mandible. *International Journal of Oral and Maxillofacial Implants* **24**(Suppl): 158–168.
- den Hartog, L., Slater, J.J., Vissink, A., Meijer, H.J. & Raghoobar, G.M. (2008) Treatment outcome of immediate, early and conventional single-tooth implants in the aesthetic zone: a systematic review to survival, bone level, soft-tissue, aesthetics and patient satisfaction. *Journal of Clinical Periodontology* **35**: 1073–1086.
- Grutter, L. & Belsler, U.C. (2009) Implant loading protocols for the partially edentulous esthetic zone. *International Journal of Oral and Maxillofacial Implants* **24**(Suppl): 169–179.
- Harder, S. & Kern, M. (2009) Survival and complications of computer aided-designing and computer-aided manufacturing vs. conventionally fabricated implant-supported reconstructions: a systematic review. *Clinical Oral Implants Research* **20**(Suppl 4): 48–54.
- Holm-Pedersen, P., Lang, N.P. & Muller, F. (2007) What are the longevities of teeth and oral implants? *Clinical Oral Implants Research* **18** (Suppl 3): 15–19.
- Iqbal, M.K. & Kim, S. (2007a) For teeth requiring endodontic treatment, what are the differences in outcomes of restored endodontically treated teeth compared to implant-supported restorations? *International Journal of Oral and Maxillofacial Implants* **22**(Suppl): 96–116.
- Jung, R.E., Pjetursson, B.E., Glauser, R., Zembic, A., Zwahlen, M. & Lang, N.P. (2008b) A systematic review of the 5-year survival and complication rates of implant-supported single crowns. *Clinical Oral Implants Research* **19**: 119–130.
- Kotsovilis, S., Fourmousis, I., Karoussis, I.K. & Bamia, C. (2009) A systematic review and meta-analysis on the effect of implant length on the survival of rough-surface dental implants. *Journal of Periodontology* **80**: 1700–1718.
- Nakamura, K., Kanno, T., Milleding, P. & Ortengren, U. (2010a) Zirconia as a dental implant abutment material: a systematic review. *International Journal of Prosthodontics* **23**: 299–309.
- Ong, C.T., Ivanovski, S., Needleman, I.G., Retzepi, M., Moles, D.R., Tonetti, M.S. & Donos, N. (2008) Systematic review of implant outcomes in treated periodontitis subjects. *Journal of Clinical Periodontology* **35**: 438–462.
- Pjetursson, B.E. & Lang, N.P. (2008) Prosthetic treatment planning on the basis of scientific evidence. *Journal of Oral Rehabilitation* **35**(Suppl 1): 72–79.
- Pjetursson, B.E., Tan, W.C., Zwahlen, M. & Lang, N.P. (2008) A systematic review of the success of sinus floor elevation and survival of implants inserted in combination with sinus floor elevation. *Journal of Clinical Periodontology* **35**: 216–240.
- Raviv, E., Turcotte, A. & Harel-Raviv, M. (2010) Short dental implants in reduced alveolar bone height. *Quintessence International* **41**: 575–579.
- Rebellato, E. & White, S.N. (2008) Readers' roundtable. Outcomes of root canal treatment and restoration, implant-supported single crowns, fixed partial dentures, and extraction without replacement: A systematic review. *Journal of Prosthetic Dentistry* **99**: 1.
- Rocuzzo, M., Aglietta, M. & Cordaro, L. (2009) Implant loading protocols for partially edentulous maxillary posterior sites. *International Journal of Oral and Maxillofacial Implants* **24**(Suppl): 147–157.
- Romeo, E., Bivio, A., Mosca, D., Scanferla, M., Ghisolfi, M. & Storelli, S. (2010a) The use of short dental implants in clinical practice: literature review. *Mineva Stomatologica* **59**: 23–31.
- Safii, S.H., Palmer, R.M. & Wilson, R.F. (2010) Risk of implant failure and marginal bone loss in subjects with a history of periodontitis: a systematic review and meta-analysis. *Clinical Implant Dentistry And Related Research* **12**: 165–174.
- Salinas, T.J. & Eckert, S.E. (2007) In patients requiring single-tooth replacement, what are the outcomes of implant- as compared to tooth-supported restorations? *International Journal of Oral and Maxillofacial Implants* **22**(Suppl): 71–95.
- Salvi, G.E. & Bragger, U. (2009) Mechanical and technical risks in implant therapy. *International Journal of Oral and Maxillofacial Implants* **24** (Suppl): 69–85.
- Schou, S., Holmstrup, P., Worthington, H.V. & Esposito, M. (2006) Outcome of implant therapy in patients with previous tooth loss due to periodontitis. *Clinical Oral Implants Research* **17** (Suppl 2): 104–123.
- Schou, S. (2008) Implant treatment in periodontitis-susceptible patients: a systematic review. *Journal of Oral Rehabilitation* **35**(Suppl 1): 9–22.
- Theoharidou, A., Petridis, H.P., Tzannas, K. & Garefis, P. (2008a) Abutment screw loosening in single-implant restorations: a systematic review. *International Journal of Oral and Maxillofacial Implants* **23**: 681–690.
- Tomasi, C., Wennstrom, J.L. & Berglundh, T. (2008) Longevity of teeth and implants - a systematic review. *Journal of Oral Rehabilitation* **35**(Suppl 1): 23–32.
- Torabinejad, M., Anderson, P., Bader, J., Brown, L.J., Chen, L.H., Goodacre, C.J., Kattadiyil, M.T., Kutsenko, D., Lozada, J., Patel, R., Petersen, F., Puterman, I. & White, S.N. (2007) Outcomes of root canal treatment and restoration, implant-supported single crowns, fixed partial dentures, and extraction without replacement: a systematic review. *Journal of Prosthetic Dentistry* **98**: 285–311.
- Torabinejad, M., Lozada, J., Puterman, I. & White, S.N. (2008) Endodontic therapy or single tooth implant? A systematic review. *Journal - California Dental Association* **36**: 429–437.
- Waasdorp, J.A., Evian, C.I. & Mandracchia, M. (2010) Immediate placement of implants into infected sites: a systematic review of the literature. *Journal of Periodontology* **81**: 801–808.
- Weber, H.P. & Sukotjo, C. (2007a) Does the type of implant prosthesis affect outcomes in the partially edentulous patient? *International Journal of Oral and Maxillofacial Implants* **22**(Suppl): 140–172.

## Included articles

- Andersson, B., Odman, P., Lindvall, A.M. & Branemark, P.I. (1998a) Five-year prospective study of prosthodontic and surgical single-tooth implant treatment in general practices and at a specialist clinic. *The International Journal of Prosthodontics* **11**: 351–355.
- Andersson, B., Odman, P., Lindvall, A.M. & Branemark, P.I. (1998b) Cemented single crowns on osseointegrated implants after 5 years: results from a prospective study on CeraOne. *The International Journal of Prosthodontics* **11**: 212–218.
- Bernard, J.P., Schatz, J.P., Christou, P., Belsler, U. & Kiliaridis, S. (2004) Long-term vertical changes of the anterior maxillary teeth adjacent to single implants in young and mature adults. A retrospective study. *Journal of Clinical Periodontology* **31**: 1024–1028.
- Bianchi, A.E. & Sanfilippo, F. (2004) Single-tooth replacement by immediate implant and connective tissue graft: a 1-9-year clinical evaluation. *Clinical Oral Implants Research* **15**: 269–277.
- Bonde, M.J., Stokholm, R., Isidor, F. & Schou, S. (2010) Outcome of implant-supported single-tooth replacements performed by dental students. A 10-year clinical and radiographic retrospective study. *European Journal of Oral Implantology* **3**: 37–46.
- Bornstein, M.M., Schmid, B., Belsler, U.C., Lussi, A. & Buser, D. (2005) Early loading of non-submerged titanium implants with a sandblasted and acid-etched surface. 5-year results of a prospective study in partially edentulous patients. *Clinical Oral Implants Research* **16**: 631–638.
- Bragger, U., Karoussis, I., Persson, R., Pjetursson, B., Salvi, G. & Lang, N. (2005) Technical and biological complications/failures with single crowns and fixed partial dentures on implants: a 10-year prospective cohort study. *Clinical Oral Implants Research* **16**: 326–334.
- De Boever, A.L. & De Boever, J.A. (2005) Guided bone regeneration around non-submerged implants in narrow alveolar ridges: a prospective long-term clinical study. *Clinical Oral Implants Research* **16**: 549–556.
- De Boever, A.L., Quirynen, M., Coucke, W., Theuniers, G. & De Boever, J.A. (2009) Clinical and radiographic study of implant treatment out-

- come in periodontally susceptible and non-susceptible patients: a prospective long-term study. *Clinical Oral Implants Research* **20**: 1341–1350.
- Degidi, M., Iezzi, G., Perrotti, V. & Piattelli, A. (2009) Comparative analysis of immediate functional loading and immediate nonfunctional loading to traditional healing periods: a 5-year follow-up of 550 dental implants. *Clinical Implant Dentistry and Related Research* **11**: 257–266.
- Elkhoury, J.S., McGlumphy, E.A., Tatakis, D.N. & Beck, F.M. (2005) Clinical parameters associated with success and failure of single-tooth titanium plasma-sprayed cylindrical implants under stricter criteria: a 5-year retrospective study. *The International Journal of Oral & Maxillofacial Implants* **20**: 687–694.
- Gibbard, L.L. & Zarb, G. (2002a) A 5-year prospective study of implant-supported single-tooth replacements. *Journal* **68**: 110–116.
- Gotfredsen, K. (2004) A 5-year prospective study of single-tooth replacements supported by the Astra Tech implant: a pilot study. *Clinical Implant Dentistry and Related Research* **6**: 1–8.
- Gotfredsen, K. (2009) A 10-Year Prospective Study of Single Tooth Implants Placed in the Anterior Maxilla. *Clinical Implant Dentistry and Related Research*. Epub August 14, 2009.
- Haas, R., Polak, C., Furhauser, R., Mailath-Pokorny, G., Dortbudak, O. & Watzek, G. (2002) A long-term follow-up of 76 Branemark single-tooth implants. *Clinical Oral Implants Research* **13**: 38–43.
- Halg, G.A., Schmid, J. & Hammerle, C.H. (2008) Bone level changes at implants supporting crowns or fixed partial dentures with or without cantilevers. *Clinical Oral Implants Research* **19**: 983–990.
- Henry, P.J., Laney, W.R., Jemt, T., Harris, D., Krogh, P.H., Polizzi, G., Zarb, G.A. & Herrmann, I. (1996) Osseointegrated implants for single-tooth replacement: a prospective 5-year multicenter study. *The International Journal of Oral & Maxillofacial Implants* **11**: 450–455.
- Jemt, T. & Lekholm, U. (2005) Single implants and buccal bone grafts in the anterior maxilla: measurements of buccal crestal contours in a 6-year prospective clinical study. *Clinical Implant Dentistry and Related Research* **7**: 127–135.
- Jemt, T. (2008b) Single implants in the anterior maxilla after 15 years of follow-up: comparison with central implants in the edentulous maxilla. *International Journal of Prosthodontics* **21**: 400–408.
- Jemt, T. (2009) Cemented CeraOne and porcelain fused to TiAdapt abutment single-implant crown restorations: a 10-year comparative follow-up study. *Clinical Implant Dentistry and Related Research* **11**: 303–310.
- Jung, R.E., Fenner, N., Hämmerle, C.H.F. & Zitzmann, N.U. (2012a) Long-term outcome of implants placed with guided bone regeneration (GBR) using resorbable and non-resorbable membranes after 12 to 14 years. *Clinical Oral Implants Research*. Epub June 16, 2012.
- Jung, R.E., Zaugg, B., Philipp, A.O.H., Truninger, T. C., Siegenthaler, D. & Hämmerle, C.H.F. (2012b) A prospective controlled clinical trial evaluating the clinical radiological and aesthetic outcome after 5 years of immediately placed implants in sockets exhibiting pericapical pathology. *Clinical Oral Implants Research*. Epub June 8, 2012.
- Kreissl, M.E., Gerds, T., Mucbe, R., Heydecke, G. & Strub, J.R. (2007) Technical complications of implant-supported fixed partial dentures in partially edentulous cases after an average observation period of 5 years. *Clinical Oral Implants Research* **18**: 720–726.
- Krennmair, G., Seemann, R., Schmidinger, S., Ewers, R. & Piehslinger, E. (2010) Clinical outcome of root-shaped dental implants of various diameters: 5-year results. *International Journal of Oral and Maxillofacial Implants* **25**: 357–366.
- Krieger, O., Matuliene, G., Husler, J., Salvi, G.E., Pietursson, B. & Bragger, U. (2009) Failures and complications in patients with birth defects restored with fixed dental prostheses and single crowns on teeth and/or implants. *Clinical Oral Implants Research* **20**: 809–816.
- Levin, L., Pathael, S., Dolev, E. & Schwartz-Arad, D. (2005) Aesthetic versus surgical success of single dental implants: 1- to 9-year follow-up. *Practical Procedures & Aesthetic Dentistry: PPAD* **17**: 533–538; quiz 540, 566.
- MacDonald, K., Pharoah, M., Todescan, R. & Deporter, D. (2009) Use of sintered porous-surfaced dental implants to restore single teeth in the maxilla: a 7- to 9-year follow-up. *International Journal of Periodontics and Restorative Dentistry* **29**: 191–199.
- Matarasso, S., Rasperini, G., Iorio Siciliano, V., Salvi, G.E., Lang, N.P. & Aglietta, M. (2010) A 10-year retrospective analysis of radiographic bone-level changes of implants supporting single-unit crowns in periodontally compromised vs. periodontally healthy patients. *Clinical Oral Implants Research* **21**: 898–903.
- Mericske-Stern, R., Grutter, L., Rosch, R. & Mericske, E. (2001) Clinical evaluation and prosthetic complications of single tooth replacements by non-submerged implants. *Clinical Oral Implants Research* **12**: 309–318.
- Palmer, R.M., Palmer, P.J. & Smith, B.J. (2000) A 5-year prospective study of Astra single tooth implants. *Clinical Oral Implants Research* **11**: 179–182.
- Pikner, S.S., Grondahl, K., Jemt, T. & Friberg, B. (2009) Marginal bone loss at implants: a retrospective, long-term follow-up of turned Branemark System implants. *Clinical Implant Dentistry and Related Research* **11**: 11–23.
- Polizzi, G., Fabbro, S., Furri, M., Herrmann, I. & Squarzone, S. (1999) Clinical application of narrow Branemark System implants for single-tooth restorations. *The International Journal of Oral & Maxillofacial Implants* **14**: 496–503.
- Romeo, E., Lops, D., Margutti, E., Ghisolfi, M., Chiapasco, M. & Vogel, G. (2004) Long-term survival and success of oral implants in the treatment of full and partial arches: a 7-year prospective study with the ITI dental implant system. *The International Journal of Oral & Maxillofacial Implants* **19**: 247–259.
- Romeo, E., Ghisolfi, M., Rozza, R., Chiapasco, M. & Lops, D. (2006) Short (8-mm) dental implants in the rehabilitation of partial and complete edentulism: a 3- to 14-year longitudinal study. *International Journal of Prosthodontics* **19**: 586–592.
- Scheller, H., Urgell, J.P., Kultje, C., Klineberg, I., Goldberg, P.V., Stevenson-Moore, P., Alonso, J.M., Schaller, M., Corria, R.M., Engquist, B., Toreskog, S., Kastenbaum, F. & Smith, C.R. (1998) A 5-year multicenter study on implant-supported single crown restorations. *The International Journal of Oral & Maxillofacial Implants* **13**: 212–218.
- Schmidlin, K., Schnell, N., Steiner, S., Salvi, G.E., Pjetursson, B., Matuliene, G., Zwahlen, M., Bragger, U. & Lang, N.P. (2010) Complication and failure rates in patients treated for chronic periodontitis and restored with single crowns on teeth and/or implants. *Clinical Oral Implants Research* **21**: 550–557.
- Schneider, D., Witt, L. & Hammerle, C.H. (2011) Influence of the crown-to-implant length ratio on the clinical performance of implants supporting single crown restorations: a cross-sectional retrospective 5-year investigation. *Clinical Oral Implants Research*. Epub June 22, 2011.
- Schropp, L. & Isidor, F. (2008) Clinical outcome and patient satisfaction following full-flap elevation for early and delayed placement of single-tooth implants: a 5-year randomized study. *International Journal of Oral and Maxillofacial Implants* **23**: 733–743.
- Taylor, R.C., McGlumphy, E.A., Tatakis, D.N. & Beck, F.M. (2004) Radiographic and clinical evaluation of single-tooth Biolok implants: a 5-year study. *The International Journal of Oral & Maxillofacial Implants* **19**: 849–854.
- Thilander, B., Odman, J. & Jemt, T. (1999) Single implants in the upper incisor region and their relationship to the adjacent teeth. An 8-year follow-up study. *Clinical Oral Implants Research* **10**: 346–355.
- Urdaneta, R.A., Rodriguez, S., McNeil, D.C., Weed, M. & Chuang, S.K. (2010) The effect of increased crown-to-implant ratio on single-tooth locking-taper implants. *International Journal of Oral and Maxillofacial Implants* **25**: 729–743.
- Vigolo, P. & Givani, A. (2000) Clinical evaluation of single-tooth mini-implant restorations: a five-year retrospective study. *The Journal of Prosthetic Dentistry* **84**: 50–54.
- Vigolo, P. & Givani, A. (2009) Platform-switched restorations on wide-diameter implants: a 5-year clinical prospective study. *International Journal of Oral and Maxillofacial Implants* **24**: 103–109.
- Wagenberg, B. & Froum, S.J. (2006) A retrospective study of 1925 consecutively placed immediate implants from 1988 to 2004. *The International Journal of Oral & Maxillofacial Implants* **21**: 71–80.
- Wennstrom, J.L., Ekstubb, A., Grondahl, K., Karlsson, S. & Lindhe, J. (2005) Implant-supported single-tooth restorations: a 5-year prospective study. *Journal of Clinical Periodontology* **32**: 567–574.
- Zafropoulos, G.G., Deli, G., Bartee, B.K. & Hoffmann, O. (2010) Single-tooth implant placement and loading in fresh and regenerated extraction sockets. Five-year results: a case series using two different implant designs. *Journal of Periodontology* **81**: 604–615.

## List of excluded full-text articles and the reason for exclusion

- Agamy, E.M. & Niedermeier, W. (2010) Indirect sinus floor elevation for osseointegrated prostheses. A 10-year prospective study. *The Journal of Oral Implantology* **36**: 113–121. [Exclusion criteria: no information on prosthetics.]
- Aglietta, M., Siciliano, V.I., Rasperini, G., Caffero, C., Lang, N.P. & Salvi, G.E. (2011) A 10-year retrospective analysis of marginal bone-level changes around implants in periodontally healthy and periodontally compromised tobacco smokers. *Clinical Oral Implants Research* **22**: 47–53. [Exclusion criteria: no information on prosthetics.]
- Andersen, E., Haanaes, H.R. & Knutsen, B.M. (2002) Immediate loading of single-tooth ITI implants in the anterior maxilla: a prospective 5-year pilot study. *Clinical Oral Implants Research* **13**: 281–287. [Exclusion criteria: < 10 patients.]
- Andreana, S., Beneduce, C. & Buhite, R. (2008) Implant success rate in dental school setting: retrospective study. *The New York State Dental Journal* **74**: 67–70. [Exclusion criteria: chart review (no clinical examination).]
- Anitua, E., Errazquin, J.M., de Pedro, J., Barrio, P., Begona, L. & Orive, G. (2010) Clinical evaluation of Tiny(R) 2.5- and 3.0-mm narrow-diameter implants as definitive implants in different clinical situations: a retrospective cohort study. *European Journal of Oral Implantology* **3**: 315–322. [Exclusion criteria: follow-up <5 years.]
- Anitua, E. & Orive, G. (2010) Short implants in maxillae and mandibles: a retrospective study with 1 to 8 years of follow-up. *Journal of Periodontology* **81**: 819–826. [Exclusion criteria: no detailed information on prosthetics.]
- Anitua, E., Orive, G., Aguirre, J.J. & Andia, I. (2008a) Five-year clinical evaluation of short dental implants placed in posterior areas: a retrospective study. *Journal of Periodontology* **79**: 42–48. [Exclusion criteria: follow-up <5 years.]
- Anitua, E., Orive, G., Aguirre, J.J. & Andia, I. (2008b) Clinical outcome of immediately loaded dental implants bioactivated with plasma rich in growth factors: a 5-year retrospective study. *Journal of Periodontology* **79**: 1168–1176. [Exclusion criteria: follow-up <5 years.]
- Anitua, E., Orive, G., Aguirre, J.J., Ardanza, B. & Andia, I. (2008c) 5-year clinical experience with BTI dental implants: risk factors for implant failure. *Journal of Clinical Periodontology* **35**: 724–732. [Exclusion criteria: follow-up <5 years.]
- Arisan, V., Bolukbasi, N., Ersanli, S. & Ozdemir, T. (2010) Evaluation of 316 narrow diameter implants followed for 5-10 years: a clinical and radiographic retrospective study. *Clinical Oral Implants Research* **21**: 296–307. [Exclusion criteria: no detailed information on prosthetics]
- Arlin, M.L. (2006) Short dental implants as a treatment option: results from an observational study in a single private practice. *The International Journal of Oral & Maxillofacial Implants* **21**: 769–776. [Exclusion criteria: follow-up <5 years.]
- Arlin, M.L. (2007) Survival and success of sand-blasted, large-grit, acid-etched and titanium plasma-sprayed implants: a retrospective study. *Journal (Canadian Dental Association)* **73**: 821. [Exclusion criteria: follow-up <5 years.]
- Avvanzo, P., Ciavarella, D., Avvanzo, A., Giannone, N., Carella, M. & Lo Muzio, L. (2009) Immediate placement and temporization of implants: three-to five-year retrospective results. *The Journal of Oral Implantology* **35**: 136–142. [Exclusion criteria: follow-up <5 years.]
- Aykent, F., Inan, O., Ozysesil, A.G. & Alptekin, N. O. (2007) A 1- to 12-year clinical evaluation of 106 endosseous implants supporting fixed and removable prostheses. *The International Journal of Periodontics & Restorative Dentistry* **27**: 358–367. [Exclusion criteria: follow-up <5 years]
- Babbush, C.A. (2006) Immediate implant placement in fresh extraction sites. *Dental Implantology Update* **17**: 89–93. [Exclusion criteria: no detailed information on prosthetics.]
- Bahat, O. (2009) Technique for placement of oxidized titanium implants in compromised maxillary bone: prospective study of 290 implants in 126 consecutive patients followed for a minimum of 3 years after loading. *The International Journal of Oral & Maxillofacial Implants* **24**: 325–334. [Exclusion criteria: follow-up <5 years.]
- Balshe, A.A., Assad, D.A., Eckert, S.E., Koka, S. & Weaver, A.L. (2009) A retrospective study of the survival of smooth- and rough-surface dental implants. *The International Journal of Oral & Maxillofacial Implants* **24**: 1113–1118. [Exclusion criteria: no detailed information on prosthetics.]
- Balshe, A.A., Eckert, S.E., Koka, S., Assad, D.A. & Weaver, A.L. (2008) The effects of smoking on the survival of smooth- and rough-surface dental implants. *The International Journal of Oral & Maxillofacial Implants* **23**: 1117–1122. [Exclusion criteria: no information on prosthetics.]
- Balshi, S.F., Wolfinger, G.J. & Balshi, T.J. (2007) A retrospective analysis of 44 implants with no rotational primary stability used for fixed prosthesis anchorage. *The International Journal of Oral & Maxillofacial Implants* **22**: 467–471. [Exclusion criteria: no detailed information on prosthetics.]
- Balshi, T.J., Wolfinger, G.J., Wulc, D. & Balshi, S.F. (2011) A prospective analysis of immediate provisionalization of single implants. *Journal of Prosthodontics: Official Journal of the American College of Prosthodontists* **20**: 10–15. [Exclusion criteria: follow-up <5 years.]
- Barone, A., Orlando, B., Tonelli, P. & Covani, U. (2011) Survival rate for implants placed in the posterior maxilla with and without sinus augmentation: a comparative cohort study. *Journal of Periodontology* **82**: 219–226. [Exclusion criteria: no detailed information on prosthetics]
- Becker, W., Goldstein, M., Becker, B.E., Sennerby, L., Kois, D. & Hujuel, P. (2009) Minimally invasive flapless implant placement: follow-up results from a multicenter study. *Journal of Periodontology* **80**: 347–352. [Exclusion criteria: follow-up <5 years.]
- Benic, G.I., Jung, R.E., Siegenthaler, D.W. & Hammerle, C.H. (2009) Clinical and radiographic comparison of implants in regenerated or native bone: 5-year results. *Clinical Oral Implants Research* **20**: 507–513. [Exclusion criteria: no detailed information on prosthetics.]
- Bersani, E., Coppede, A.R. & de Paula Pinto Prata, H.H. (2010) Immediate loading of implants placed in fresh extraction sockets in the molar area with flapless and graftless procedures: a case series. *The International Journal of Periodontics & Restorative Dentistry* **30**: 291–299. [Exclusion criteria: no detailed information on prosthetics.]
- Binahmed, A., Stoykewych, A., Hussain, A., Love, B. & Pruthi, V. (2007) Long-term follow-up of hydroxyapatite-coated dental implants—a clinical trial. *The International Journal of Oral & Maxillofacial Implants* **22**: 963–968. [Exclusion criteria: no detailed information on prosthetics]
- Bornstein, M.M., Chappuis, V., von Arx, T. & Buser, D. (2008) Performance of dental implants after staged sinus floor elevation procedures: 5-year results of a prospective study in partially edentulous patients. *Clinical Oral Implants Research* **19**: 1034–1043. [Exclusion criteria: no detailed information on prosthetics.]
- Brennan, M., Houston, F., O'Sullivan, M. & O'Connell, B. (2010) Demographics of implant placement and complications of a patient subgroup in a dental hospital population. *Journal of the Irish Dental Association* **56**: 85–92. [Exclusion criteria: no detailed information on prosthetics.]
- Buser, D., Dula, K., Lang, N.P. & Nyman, S. (1996) Long-term stability of osseointegrated implants in bone regenerated with the membrane technique. 5-year results of a prospective study with 12 implants. *Clinical Oral Implants Research* **7**: 175–183. [Exclusion criteria: < 10 patients.]
- Carbone, M., Goss, E., Borione, M., Bava, L., Broccoletti, R., Carrozzo, M. & Gandolfo, S. (2007) Implant supported prostheses with Bone System implant system: a retrospective study with follow-up period up to 13-years about 1021 fixtures. *Minerva Stomatologica* **56**: 481–495. [Exclusion criteria: no detailed information on prosthetics.]
- Cardemil, C., Ristevski, Z., Alsen, B. & Dahlin, C. (2009) Influence of different operator setups on implant survival rate: a retrospective clinical study. *Clinical Implant Dentistry and Related Research* **11**: 288–291.
- Carinci, F., Brunelli, G. & Danza, M. (2009a) Platform switching and bone platform switching. *The Journal of Oral Implantology* **35**: 245–250. [Exclusion criteria: no detailed information on prosthetics.]
- Carinci, F., Brunelli, G., Zollino, I., Franco, M., Viscioni, A., Rigo, L., Guidi, R. & Strohmer, L. (2009b) Mandibles grafted with fresh-frozen bone: an evaluation of implant outcome. *Implant Dentistry* **18**: 86–95. [Exclusion criteria: no detailed information on prosthetics.]
- Caubet, J., Petzold, C., Saez-Torres, C., Morey, M., Iriarte, J.I., Sanchez, J., Torres, J.J., Ramis, J.M. & Monjo, M. (2011) Sinus graft with safescraper: 5-year results. *Journal of Oral and Maxillofacial Surgery* **69**: 482–490. [Exclusion criteria: no detailed information on prosthetics.]



- Cavalcanti, R., Oreglia, F., Manfredonia, M.F., Gianserra, R. & Esposito, M. (2011) The influence of smoking on the survival of dental implants: a 5-year pragmatic multicentre retrospective cohort study of 1727 patients. *European Journal of Oral Implantology* 4: 39–45. [Exclusion criteria: no detailed information on prosthetics.]
- Chiapasco, M. (2004) Early and immediate restoration and loading of implants in completely edentulous patients. *The International Journal of Oral & Maxillofacial Implants* 19(Suppl): 76–91.
- Christensen, G.J. (2008) Three-unit fixed prostheses versus implant-supported single crowns. *Journal of the American Dental Association* 139: 191–194. [Exclusion criteria: no clinical study.]
- Chung, W.E., Rubenstein, J.E., Phillips, K.M. & Raigrodski, A.J. (2009) Outcomes assessment of patients treated with osseointegrated dental implants at the University of Washington Graduate Prosthodontic Program, 1988 to 2000. *The International Journal of Oral & Maxillofacial Implants* 24: 927–935. [Exclusion criteria: not all patients clinically examined.]
- Cicciu, M., Beretta, M., Risitano, G. & Maiorana, C. (2008) Cemented-retained vs screw-retained implant restorations: an investigation on 1939 dental implants. *Minerva Stomatologica* 57: 167–179. [Exclusion criteria: follow-up <5 years.]
- Cochran, D., Oates, T., Morton, D., Jones, A., Buser, D. & Peters, F. (2007) Clinical field trial examining an implant with a sand-blasted, acid-etched surface. *Journal of Periodontology* 78: 974–982. [Exclusion criteria: no information on prosthetics.]
- Conrad, H.J., Jung, J., Barczak, M., Basu, S. & Seong, W.J. (2011) Retrospective cohort study of the predictors of implant failure in the posterior maxilla. *The International Journal of Oral & Maxillofacial Implants* 26: 154–162. [Exclusion criteria: no detailed information on prosthetics.]
- Cordaro, L., Torsello, F. & Rocuzzo, M. (2009b) Implant loading protocols for the partially edentulous posterior mandible. *The International Journal of Oral & Maxillofacial Implants* 24(Suppl): 158–168.
- Corinaldesi, G., Pieri, F., Sapigni, L. & Marchetti, C. (2009) Evaluation of survival and success rates of dental implants placed at the time of or after alveolar ridge augmentation with an autogenous mandibular bone graft and titanium mesh: a 3- to 8-year retrospective study. *The International Journal of Oral & Maxillofacial Implants* 24: 1119–1128. [Exclusion criteria: no detailed information on prosthetics.]
- Cosyn, J. & De Rouck, T. (2009) Aesthetic outcome of single-tooth implant restorations following early implant placement and guided bone regeneration: crown and soft tissue dimensions compared with contralateral teeth. *Clinical Oral Implants Research* 20: 1063–1069. [Exclusion criteria: follow-up <5 years.]
- Creton, M., Cune, M., Verhoeven, W., Muradin, M., Wismeijer, D. & Meijer, G. (2010) Implant treatment in patients with severe hypodontia: a retrospective evaluation. *Journal of Oral and Maxillofacial Surgery* 68: 530–538. [Exclusion criteria: no detailed information on prosthetics]
- Cune, M.S., Strooker, H., van der Reijden, W.A., de Putter, C., Laine, M.L. & Verhoeven, J.W. (2009) Dental implants in persons with severe epilepsy and multiple disabilities: a long-term retrospective study. *The International Journal of Oral & Maxillofacial Implants* 24: 534–540. [Exclusion criteria: no information on prosthetics.]
- Danza, M., Riccardo, G. & Carinci, F. (2010) Bone platform switching: a retrospective study on the slope of reverse conical neck. *Quintessence International* 41: 35–40. [Exclusion criteria: no detailed information on prosthetics.]
- Degidi, M., Nardi, D. & Piattelli, A. (2008a) Peri-implant tissue and radiographic bone levels in the immediately restored single-tooth implant: a retrospective analysis. *Journal of Periodontology* 79: 252–259. [Exclusion criteria: follow-up < 5 years.]
- Degidi, M., Piattelli, A. & Carinci, F. (2008b) Clinical outcome of narrow diameter implants: a retrospective study of 510 implants. *Journal of Periodontology* 79: 49–54. [Exclusion criteria: no information on prosthetics.]
- Degidi, M., Piattelli, A., Iezzi, G. & Carinci, F. (2007a) Immediately loaded short implants: analysis of a case series of 133 implants. *Quintessence International* 38: 193–201. [Exclusion criteria: follow-up < 5 years.]
- Degidi, M., Piattelli, A., Iezzi, G. & Carinci, F. (2007b) Do longer implants improve clinical outcome in immediate loading? *Journal Of Oral And Maxillofacial Surgery: Official Journal of the American Association of Oral And Maxillofacial Surgeons* 36: 1172–1176. [Exclusion criteria: follow-up < 5 years.]
- Degidi, M., Piattelli, A., Iezzi, G. & Carinci, F. (2007c) Wide-diameter implants: analysis of clinical outcome of 304 fixtures. *Journal of Periodontology* 78: 52–58. [Exclusion criteria: follow-up < 5 years.]
- Deporter, D., Ogiso, B., Sohn, D.S., Ruljancich, K. & Pharoah, M. (2008) Ultrashort sintered porous-surfaced dental implants used to replace posterior teeth. *Journal of Periodontology* 79: 1280–1286. [Exclusion criteria: < 10 patients with SCI.]
- Di Felice, R., D'Amario, M., De Dominicis, A., Garocchio, S., D'Arcangelo, C. & Giannoni, M. (2011) Immediate placement of bone level Straumann implants: a case series. *The International Journal of Periodontics & Restorative Dentistry* 31: 57–65. [Exclusion criteria: < 10 patients with SCI.]
- Doyle, S.L., Hodges, J.S., Pesun, I.J., Law, A.S. & Bowles, W.R. (2006) Retrospective cross sectional comparison of initial nonsurgical endodontic treatment and single-tooth implants. *Journal of Endodontics* 32: 822–827. [Exclusion criteria: chart review (no clinical examination).]
- Duminil, G., Muller-Bolla, M., Brun, J.P., Leclercq, P., Bernard, J.P. & Dohan Ehrenfest, D.M. (2008) Success rate of the EVL evolution implants (SERF): a five-year longitudinal multicenter study. *The Journal of Oral Implantology* 34: 282–289. [Exclusion criteria: no detailed information on prosthetics.]
- Farino, M., Branscum, A., Robinson, F.G., Jasper, S., Al-Sabbagh Dds, M., Puleo, D.A. & Thomas, M.V. (2010) Programmatic effectiveness of a university-based implant training program: long-term, patient-centered outcomes. *Journal of Long-Term Effects of Medical Implants* 20: 343–351. [Exclusion criteria: no detailed information on prosthetics.]
- Franchini, I., Capelli, M., Fumagalli, L., Parenti, A. & Testori, T. (2011) Multicenter retrospective analysis of 201 consecutively placed camlog dental implants. *The International Journal of Periodontics & Restorative Dentistry* 31: 255–263. [Exclusion criteria: no detailed information on prosthetics.]
- Franco, M., Rigo, L., Viscione, A., De Santis, B., Tropina, E., Brunelli, G., Guidi, R., Avantaggiato, A. & Carinci, F. (2009a) CaPO4 blasted implants inserted into iliac crest homologue frozen grafts. *The Journal of Oral Implantology* 35: 176–180. [Exclusion criteria: no detailed information on prosthetics.]
- Franco, M., Viscioni, A., Rigo, L., Guidi, R., Zollino, I., Avantaggiato, A. & Carinci, F. (2009b) Clinical outcome of narrow diameter implants inserted into allografts. *Journal of Applied Oral Science: Revista Fob* 17: 301–306. [Exclusion criteria: no detailed information on prosthetics.]
- Friberg, B. & Jemt, T. (2010) Clinical experience of TiUnite implants: a 5-year cross-sectional, retrospective follow-up study. *Clinical implant dentistry and related research* 12(Suppl 1): e95–103. [Exclusion criteria: no detailed information on prosthetics.]
- Froum, S.J., Cho, S.C., Cho, Y.S., Elian, N. & Tarnow, D. (2007) Narrow-diameter implants: a restorative option for limited interdental space. *The International Journal of Periodontics & Restorative Dentistry* 27: 449–455. [Exclusion criteria: follow-up < 5 years.]
- Fugazzotto, P.A. (2008a) Implant placement at the time of maxillary molar extraction: treatment protocols and report of results. *Journal of Periodontology* 79: 216–223. [Exclusion criteria: follow-up < 5 years]
- Fugazzotto, P.A. (2008b) Implant placement at the time of mandibular molar extraction: description of technique and preliminary results of 341 cases. *Journal of Periodontology* 79: 737–747. [Exclusion criteria: no detailed information on prosthetics.]
- Garlini, G., Redemagni, M., Donini, M. & Maiorana, C. (2010) Maxillary sinus elevation with an alloplastic material and implants: 11 years of clinical and radiologic follow-up. *Journal of Oral and Maxillofacial Surgery* 68: 1152–1157. [Exclusion criteria: no detailed information on prosthetics.]
- Gatti, C., Gatti, F., Chiapasco, M. & Esposito, M. (2008) Outcome of dental implants in partially edentulous patients with and without a history of periodontitis: a 5-year interim analysis of a cohort study. *European Journal of Oral Implantology* 1: 45–51. [Exclusion criteria: no detailed information on prosthetics.]
- George, K.M., Choi, Y.G., Rieck, K.L., Van Ess, J., Ivancakova, R. & Carr, A.B. (2011) Immediate restoration with ti-unite implants: practice-based evidence compared with animal study outcomes. *The International Journal of Prosthodontics* 24: 199–203. [Exclusion criteria: chart review (no clinical examination).]

- Gianserra, R., Cavalcanti, R., Oreglia, F., Manfredonia, M.F. & Esposito, M. (2010) Outcome of dental implants in patients with and without a history of periodontitis: a 5-year pragmatic multicentre retrospective cohort study of 1727 patients. *European Journal of Oral Implantology* **3**: 307–314. [Exclusion criteria: no detailed information on prosthetics.]
- Gibbard, L.L. & Zarb, G. (2002b) A 5-year prospective study of implant-supported single-tooth replacements. *Journal (Canadian Dental Association)* **68**: 110–116.
- Grant, B.T., Pancko, F.X. & Kraut, R.A. (2009) Outcomes of placing short dental implants in the posterior mandible: a retrospective study of 124 cases. *Journal of Oral and Maxillofacial Surgery* **67**: 713–717. [Exclusion criteria: follow-up < 5 years.]
- Grossmann, Y. & Levin, L. (2007) Success and survival of single dental implants placed in sites of previously failed implants. *Journal of Periodontology* **78**: 1670–1674. [Exclusion criteria: follow-up < 5 years.]
- Holahan, C.M., Koka, S., Kennel, K.A., Weaver, A.L., Assad, D.A., Regennitter, F.J. & Kademani, D. (2008) Effect of osteoporotic status on the survival of titanium dental implants. *The International Journal of Oral & Maxillofacial Implants* **23**: 905–910. [Exclusion criteria: no information on prosthetics.]
- Hu, X.L., Li, J.H., Luo, J., Qiu, L.X. & Lin, Y. (2011) Multidisciplinary management of congenitally missing teeth with osseointegrated dental implants: a long-term report. *The Chinese journal of dental research: the official journal of the Scientific Section of the Chinese Stomatological* **14**: 29–36. [Exclusion criteria: < 10 patients with SCL.]
- Hussaini, S., Weiner, S. & Ahmad, M. (2010) Implant survival rates in a condensed surgical and prosthetic training program for general practitioners in dental implants. *Implant Dentistry* **19**: 73–80. [Exclusion criteria: no information on prosthetics.]
- Iqbal, M.K. & Kim, S. (2007b) For teeth requiring endodontic treatment, what are the differences in outcomes of restored endodontically treated teeth compared to implant-supported restorations? *The International Journal of Oral & Maxillofacial Implants* **22**(Suppl): 96–116.
- Jebreen, S.E. & Khraisat, A. (2007) Multicenter retrospective study of ITI implant-supported posterior partial prosthesis in Jordan. *Clinical implant dentistry and related research* **9**: 89–93. [Exclusion criteria: no detailed information on prosthetics.]
- Jemt, T., Ahlberg, G., Henriksson, K. & Bondevik, O. (2006) Changes of anterior clinical crown height in patients provided with single-implant restorations after more than 15 years of follow-up. *The International Journal of Prosthodontics* **19**: 455–461. [Exclusion criteria: multiple publications on same patient cohort.]
- Jemt, T., Ahlberg, G., Henriksson, K. & Bondevik, O. (2007) Tooth movements adjacent to single-implant restorations after more than 15 years of follow-up. *The International Journal of Prosthodontics* **20**: 626–632. [Exclusion criteria: multiple publications on same patient cohort.]
- Jung, U.W., Choi, J.Y., Kim, C.S., Cho, K.S., Chai, J.K., Kim, C.K. & Choi, S.H. (2008c) Evaluation of mandibular posterior single implants with two different surfaces: a 5-year comparative study. *Journal of Periodontology* **79**: 1857–1863. [Exclusion criteria: no information on prosthetics.]
- Juodzbalsys, G., Raustia, A.M. & Kubilius, R. (2007) A 5-year follow-up study on one-stage implants inserted concomitantly with localized alveolar ridge augmentation. *Journal of Oral Rehabilitation* **34**: 781–789. [Exclusion criteria: no information on prosthetics.]
- Kacer, C.M., Dyer, J.D. & Kraut, R.A. (2010) Immediate loading of dental implants in the anterior and posterior mandible: a retrospective study of 120 cases. *Journal of Oral and Maxillofacial Surgery* **68**: 2861–2867. [Exclusion criteria: no detailed information on prosthetics]
- Kallus, T., Bessing, C., Homsy, G. & Eklund, I. (2009) Five-year evaluation of Lifecore Restore implants: a retrospective comparison with Nobel Biocare MK II implants. *Clinical implant dentistry and related research* **11**: 167–177. [Exclusion criteria: no detailed information on prosthetics]
- Kim, Y.K., Kim, S.G., Oh, H.K., Choi, Y.G., Cho, Y. S., Oh, Y.H., Son, J.S. & Ong, J.L. (2009) Evaluation of peri-implant tissue in nonsubmerged dental implants: a multicenter retrospective study. *Oral Surgery, Oral Medicine, Oral Pathology, Oral Radiology, And Endodontics* **108**: 189–195. [Exclusion criteria: follow-up < 5 years.]
- Kim, Y.K., Kim, S.G., Yun, P.Y., Hwang, J.W. & Son, M.K. (2010) Prognosis of single molar implants: a retrospective study. *The International Journal of Periodontics & Restorative Dentistry* **30**: 401–407. [Exclusion criteria: follow-up < 5 years.]
- Knauf, M., Gerds, T., Mucbe, R. & Strub, J.R. (2007) Survival and success rates of 3i implants in partially edentulous patients: results of a prospective study with up to 84-months' follow-up. *Quintessence International* **38**: 643–651. [Exclusion criteria: no information on prosthetics.]
- Koldsland, O.C., Scheie, A.A. & Aass, A.M. (2009) Prevalence of implant loss and the influence of associated factors. *Journal of Periodontology* **80**: 1069–1075. [Exclusion criteria: no detailed information on prosthetics.]
- Koo, K.T., Wikesjo, U.M., Park, J.Y., Kim, T.I., Seol, Y.J., Ku, Y., Rhyu, I.C., Chung, C.P. & Lee, Y.M. (2010) Evaluation of single-tooth implants in the second molar region: a 5-year life-table analysis of a retrospective study. *Journal of Periodontology* **81**: 1242–1249. [Exclusion criteria: no information on prosthetics.]
- Kronstrom, M., McGrath, L. & Chaytor, D. (2008) Implant dentistry in the undergraduate dental education program at Dalhousie University. Part 1: clinical outcomes. *The International Journal of Prosthodontics* **21**: 124–128. [Exclusion criteria: no detailed information on prosthetics.]
- Laviv, A., Levin, L., Usiel, Y. & Schwartz-Arad, D. (2010) Survival of immediately provisionalized dental implants: a case-control study with up to 5 years follow-up. *Clinical implant dentistry and related research* **12**(Suppl 1): e23–27. [Exclusion criteria: no detailed information on prosthetics.]
- Levin, B.P. (2011) Immediate temporization of immediate implants in the esthetic zone: evaluating survival and bone maintenance. *Compendium of Continuing Education in Dentistry* **32**: 52–5658–60, 62. [Exclusion criteria: follow-up < 5 years.]
- Levin, L., Hertzberg, R., Har-Nes, S. & Schwartz-Arad, D. (2008) Long-term marginal bone loss around single dental implants affected by current and past smoking habits. *Implant Dentistry* **17**: 422–429. [Exclusion criteria: no information on prosthetics.]
- Levin, L., Laviv, A. & Schwartz-Arad, D. (2006a) Long-term success of implants replacing a single molar. *Journal of Periodontology* **77**: 1528–1532. [Exclusion criteria: follow-up < 5 years.]
- Levin, L., Nitzan, D. & Schwartz-Arad, D. (2007) Success of dental implants placed in intraoral block bone grafts. *Journal of Periodontology* **78**: 18–21. [Exclusion criteria: no information on prosthetics.]
- Levin, L., Sadet, P. & Grossmann, Y. (2006b) A retrospective evaluation of 1,387 single-tooth implants: a 6-year follow-up. *Journal of Periodontology* **77**: 2080–2083. [Exclusion criteria: follow-up < 5 years.]
- Machtei, E.E., Mahler, D., Oettinger-Barak, O., Zuabi, O. & Horwitz, J. (2008) Dental implants placed in previously failed sites: survival rate and factors affecting the outcome. *Clinical Oral Implants Research* **19**: 259–264. [Exclusion criteria: no information on prosthetics.]
- Mahn, D.H. & Polack, M.A. (2008) Replacement of maxillary central incisors with implants. *Dentistry Today* **27**, **160**, **162**: 164. [Exclusion criteria: < 10 patients with SCL.]
- Malo, P. & de Araujo Nobre, M. (2011) Implants (3.3 mm diameter) for the rehabilitation of edentulous posterior regions: a retrospective clinical study with up to 11 years of follow-up. *Clinical implant dentistry and related research* **13**: 95–103. [Exclusion criteria: no information on prosthetics.]
- Malo, P., de Araujo Nobre, M. & Rangert, B. (2007a) Short implants placed one-stage in maxillae and mandibles: a retrospective clinical study with 1 to 9 years of follow-up. *Clinical implant dentistry and related research* **9**: 15–21. [Exclusion criteria: no information on prosthetics.]
- Malo, P., de Araujo Nobre, M. & Rangert, B. (2007b) Implants placed in immediate function in periodontally compromised sites: a five-year retrospective and one-year prospective study. *The Journal of Prosthetic Dentistry* **97**: S86–95. [Exclusion criteria: no information on prosthetics.]
- Mangano, C., Mangano, F., Shibli, J.A., Tettamanti, L., Figliuzzi, M., d'Avila, S., Sammons, R.L. & Piattelli, A. (2011) Prospective evaluation of 2,549 Morse taper connection implants: 1- to 6-year data. *Journal of Periodontology* **82**: 52–61. [Exclusion criteria: no detailed information on prosthetics.]
- Manor, Y., Oubaid, S., Mardinger, O., Chaushu, G. & Nissan, J. (2009) Characteristics of early versus late implant failure: a retrospective study. *Journal of Oral and Maxillofacial Surgery* **67**: 2649–2652. [Exclusion criteria: no detailed information on prosthetics.]
- Manso, M.C. & Wassal, T. (2010) A 10-year longitudinal study of 160 implants simultaneously installed in severely atrophic posterior maxillas grafted with autogenous bone and a synthetic bioactive resorbable graft. *Implant Dentistry* **19**:

- 351–360. [Exclusion criteria: no information on prosthetics.]
- Matsui, Y., Ohno, K., Nishimura, A., Shirota, T., Kim, S. & Miyashita, H. (2007) Long-term study of dental implants placed into alveolar cleft sites. *The Cleft palate-craniofacial journal: official publication of the American Cleft Palate-Craniofacial Association* **44**: 444–447. [Exclusion criteria: no information on prosthetics.]
- Mengel, R., Behle, M. & Flores-de-Jacoby, L. (2007) Osseointegrated implants in subjects treated for generalized aggressive periodontitis: 10-year results of a prospective, long-term cohort study. *Journal of Periodontology* **78**: 2229–2237. [Exclusion criteria: no detailed information on prosthetics.]
- Mijiritsky, E., Mardinger, O., Mazor, Z. & Chaushu, G. (2009) Immediate provisionalization of single-tooth implants in fresh-extraction sites at the maxillary esthetic zone: up to 6 years of follow-up. *Implant Dentistry* **18**: 326–333. [Exclusion criteria: provisionalization/no definitive crowns and no information.]
- Misch, C.E., Misch-Dietsh, F., Silc, J., Barboza, E., Cianciola, L.J. & Kazor, C. (2008) Posterior implant single-tooth replacement and status of adjacent teeth during a 10-year period: a retrospective report. *Journal of Periodontology* **79**: 2378–2382. [Exclusion criteria: follow-up < 5 years.]
- Misch, C.E., Steingga, J., Barboza, E., Misch-Dietsh, F., Cianciola, L.J. & Kazor, C. (2006) Short dental implants in posterior partial edentulism: a multicenter retrospective 6-year case series study. *Journal of Periodontology* **77**: 1340–1347. [Exclusion criteria: no detailed information on prosthetics.]
- Molly, L., Quirynen, M., Michiels, K. & van Steenberghe, D. (2006) Comparison between jaw bone augmentation by means of a stiff occlusive titanium membrane or an autologous hip graft: a retrospective clinical assessment. *Clinical Oral Implants Research* **17**: 481–487. [Exclusion criteria: no detailed information on prosthetics.]
- Nakamura, K., Kanno, T., Milleding, P. & Ortengren, U. (2010b) Zirconia as a dental implant abutment material: a systematic review. *The International Journal of Prosthodontics* **23**: 299–309.
- Nedir, R., Nurdin, N., Vazquez, L., Szmukler-Moncler, S., Bischof, M. & Bernard, J.P. (2010) Osteotome sinus floor elevation technique without grafting: a 5-year prospective study. *Journal of Clinical Periodontology* **37**: 1023–1028. [Exclusion criteria: < 10 patients with SCI.]
- Nelson, K., Semper, W., Hildebrand, D. & Ozyuvaci, H. (2008) A retrospective analysis of sandblasted, acid-etched implants with reduced healing times with an observation period of up to 5 years. *The International Journal of Oral & Maxillofacial Implants* **23**: 726–732. [Exclusion criteria: no detailed information on prosthetics.]
- Nixon, K.C., Chen, S.T. & Ivanovski, S. (2009) A retrospective analysis of 1,000 consecutively placed implants in private practice. *Australian Dental Journal* **54**: 123–129. [Exclusion criteria: no detailed information on prosthetics.]
- Oliva, J., Oliva, X. & Oliva, J.D. (2010) Five-year success rate of 831 consecutively placed Zirconia dental implants in humans: a comparison of three different rough surfaces. *The International Journal of Oral & Maxillofacial Implants* **25**: 336–344. [Exclusion criteria: no detailed information on prosthetics.]
- Rocuzzo, M., De Angelis, N., Bonino, L. & Aglietta, M. (2010) Ten-year results of a three-arm prospective cohort study on implants in periodontally compromised patients. Part 1: implant loss and radiographic bone loss. *Clinical Oral Implants Research* **21**: 490–496. [Exclusion criteria: no detailed information on prosthetics.]
- Romanos, G.E. & Nentwig, G.H. (2009) Immediate functional loading in the maxilla using implants with platform switching: five-year results. *The International Journal of Oral & Maxillofacial Implants* **24**: 1106–1112. [Exclusion criteria: edentulous patients/no single crowns.]
- Romeo, E., Bivio, A., Mosca, D., Scanferla, M., Ghisolfi, M. & Storelli, S. (2010b) The use of short dental implants in clinical practice: literature review. *Minerva Stomatologica* **59**: 23–31.
- Roos-Jansaker, A.M. (2007) Long time follow up of implant therapy and treatment of peri-implantitis. *Swedish Dental Journal. Supplement* **188**: 7–66. [Exclusion criteria: no information on prosthetics.]
- Sanchez-Perez, A., Moya-Villaescusa, M.J. & Caffesse, R.G. (2007) Tobacco as a risk factor for survival of dental implants. *Journal of Periodontology* **78**: 351–359. [Exclusion criteria: no detailed information on prosthetics.]
- Scarano, A., Piattelli, A., Assenza, B., Quaranta, A., Perrotti, V., Piattelli, M. & Iezzi, G. (2010) Porcine bone used in sinus augmentation procedures: a 5-year retrospective clinical evaluation. *Journal of Oral and Maxillofacial Surgery* **68**: 1869–1873. [Exclusion criteria: mixed data, no information on single crowns.]
- Schlegel, K.A., Karascholi, T., Fenner, M. & Nkenke, E. (2007) [Clinical and radiological results after augmentation procedures - a prospective study]. *Mund-, Kiefer- Und Gesichtschirurgie: Mkg* **11**: 209–219. [Exclusion criteria: mixed data, no information on single crowns.]
- Schulte, J., Flores, A.M. & Weed, M. (2007) Crown-to-implant ratios of single tooth implant-supported restorations. *The Journal of Prosthetic Dentistry* **98**: 1–5. [Exclusion criteria: chart review (no clinical examination).]
- Schwartz-Arad, D., Laviv, A. & Levin, L. (2008) Failure causes, timing, and cluster behavior: an 8-year study of dental implants. *Implant Dentistry* **17**: 200–207. [Exclusion criteria: chart review (no clinical examination).]
- Shabestari, G.O., Shayesteh, Y.S., Khojasteh, A., Alikhasi, M., Moslemi, N., Aminian, A., Masaeli, R., Eslami, B. & Treister, N.S. (2010) Implant placement in patients with oral bisphosphonate therapy: a case series. *Clinical implant dentistry and related research* **12**: 175–180. [Exclusion criteria: no detailed information on prosthetics.]
- Simonis, P., Dufour, T. & Tenenbaum, H. (2010) Long-term implant survival and success: a 10-16-year follow-up of non-submerged dental implants. *Clinical Oral Implants Research* **21**: 772–777. [Exclusion criteria: no detailed information on failed implants (SCI).]
- Smith, L.P., Ng, M., Grubor, D. & Chandu, A. (2009) Outcomes of dental implants placed in a surgical training programme. *Australian Dental Journal* **54**: 361–367. [Exclusion criteria: no detailed information on prosthetics.]
- Sohn, D.S., Kim, W.S., Lee, W.H., Jung, H.S. & Shin, I.H. (2010) A retrospective study of sintered porous-surfaced dental implants in restoring the edentulous posterior mandible: up to 9 years of functioning. *Implant Dentistry* **19**: 409–418. [Exclusion criteria: follow-up < 5 years.]
- Takahashi, T., Inai, T., Kochi, S., Fukuda, M., Yamaguchi, T., Matsui, K., Echigo, S. & Watanabe, M. (2008) Long-term follow-up of dental implants placed in a grafted alveolar cleft: evaluation of alveolar bone height. *Oral Surgery, Oral Medicine, Oral Pathology, Oral Radiology, And Endodontics* **105**: 297–302. [Exclusion criteria: no detailed information on prosthetics.]
- Tetsch, J., Tetsch, P. & Lysek, D.A. (2010) Long-term results after lateral and osteotome technique sinus floor elevation: a retrospective analysis of 2190 implants over a time period of 15 years. *Clinical Oral Implants Research* **21**: 497–503. [Exclusion criteria: no detailed information on prosthetics.]
- Theoharidou, A., Petridis, H.P., Tzannas, K. & Garefis, P. (2008b) Abutment screw loosening in single-implant restorations: a systematic review. *The International Journal of Oral & Maxillofacial Implants* **23**: 681–690.
- Tseng, C.C., Pang, I.C., Wen, M.J., Huang, K.C. & Chang, J.C. (2009) Clinical effectiveness of International Team for Oral Implantology dental implant treatment in Taiwan: a seven-year longitudinal study. *Implant Dentistry* **18**: 67–74. [Exclusion criteria: no detailed information on prosthetics.]
- Urban, I.A., Jovanovic, S.A. & Lozada, J.L. (2009) Vertical ridge augmentation using guided bone regeneration (GBR) in three clinical scenarios prior to implant placement: a retrospective study of 35 patients 12 to 72 months after loading. *The International Journal of Oral & Maxillofacial Implants* **24**: 502–510. [Exclusion criteria: no detailed information on prosthetics.]
- Urban, I.A. & Lozada, J.L. (2010) Implants placed in augmented sinuses with minimal and moderate residual crestal bone: results after 1 to 5 years. *The International Journal of Oral & Maxillofacial Implants* **25**: 1203–1212. [Exclusion criteria: no detailed information on prosthetics.]
- Viscioni, A., Franco, M., Rigo, L., Guidi, R., Brunelli, G. & Carinci, F. (2009a) Implants inserted into homografts bearing fixed restorations. *The International Journal of Prosthodontics* **22**: 148–154. [Exclusion criteria: no detailed information on prosthetics.]
- Viscioni, A., Franco, M., Rigo, L., Guidi, R., Spinelli, G. & Carinci, F. (2009b) Retrospective study of standard-diameter implants inserted into allografts. *Journal of Oral and Maxillofacial Surgery* **67**: 387–393. [Exclusion criteria: no detailed information on prosthetics.]
- Viscioni, A., Rigo, L., Franco, M., Brunelli, G., Avantaggiato, A., Sollazzo, V. & Carinci, F. (2010) Reconstruction of severely atrophic jaws using

- homografts and simultaneous implant placement: a retrospective study. *The Journal of Oral Implantology* **36**: 131–139. [Exclusion criteria: no detailed information on prosthetics.]
- Visser, A., Raghoobar, G.M., Meijer, H.J., Meijndert, L. & Vissink, A. (2011) Care and aftercare related to implant-retained dental crowns in the maxillary aesthetic region: a 5-year prospective randomized clinical trial. *Clinical Implant Dentistry and Related Research* **13**: 157–167. [Exclusion criteria: follow-up < 5 years.]
- Wagenberg, B. & Froum, S.J. (2010) Prospective study of 94 platform-switched implants observed from 1992 to 2006. *The International Journal of Periodontics & Restorative Dentistry* **30**: 9–17. [Exclusion criteria: no detailed information on prosthetics.]
- Walton, T.R. (2009a) Changes in patient and FDP profiles following the introduction of osseointegrated implant dentistry in a prosthodontic practice. *The International Journal of Prosthodontics* **22**: 127–135. [Exclusion criteria: only demographic data.]
- Walton, T.R. (2009b) Changes in the outcome of metal-ceramic tooth-supported single crowns and FDPs following the introduction of osseointegrated implant dentistry into a prosthodontic practice. *The International Journal of Prosthodontics* **22**: 260–267. [Exclusion criteria: reports on tooth-supported crowns only.]
- Weber, H.P. & Sukotjo, C. (2007b) Does the type of implant prosthesis affect outcomes in the partially edentulous patient? *The International Journal of Oral & Maxillofacial Implants* **22**(Suppl): 140–172.
- Zachrisson, B.U. (2006) Single implant-supported crowns in the anterior maxilla—potential esthetic long-term (> 5 years) problems. *World Journal of Orthodontics* **7**: 306–312. [Exclusion criteria: no detailed information on prosthetics.]